



THE POLAR WILDLIFE REPORT

Susan Crockford



The Polar Wildlife Report

Susan Crockford

Report 63, The Global Warming Policy Foundation

© Copyright 2023, The Global Warming Policy Foundation



Contents

About the author	iii
Foreword	v
Executive summary	v
1. Introduction	1
2. The Arctic	1
Sea ice and primary productivity	1
Polar bears	3
Walrus, seals, and whales	5
Birds and fish	8
3. The Antarctic	9
Sea ice and primary productivity	9
Whales and seals	10
Penguins	13
Other birds and fish	15
4. Discussion	15
Bibliography	18
References	32
Review process	36
About the Global Warming Policy Foundation	36

About the author

Dr Susan Crockford is an evolutionary biologist and has been working for more than 40 years in archaeozoology, paleozoology and forensic zoology.¹ She is a former adjunct professor at the University of Victoria, British Columbia and works full time for a private consulting company she co-owns (Pacific Identifications Inc). She is the author of *Rhythms of Life: Thyroid Hormone and the Origin of Species*, *Eaten: A Novel* (a polar bear attack thriller), *Polar Bear Facts and Myths* (for ages seven and up, also available in French, German, Dutch, Portuguese, and Norwegian), *Polar Bears Have Big Feet* (for preschoolers), and the fully referenced *How the Polar Bear Came to Be* (upcoming in 2023), *Sir David Attenborough and the Walrus Deception*, *The Polar Bear Catastrophe That Never Happened* and *Polar Bears: Outstanding Survivors of Climate Change*,² as well as a scientific paper on polar bear conservation status and a peer-reviewed paper on the distribution of ancient polar bear remains.³ She has authored several earlier briefing papers, reports, and videos for GWPF, as well as opinion pieces for major news outlets, on polar bear and walrus ecology and conservation.⁴ Susan Crockford blogs at www.polarbearsience.com.



Foreword

This *Polar Wildlife Report* is intended to provide a brief view of the habitat and conservation status of critical wildlife species in the Arctic and Antarctic, with commentary on inconsistencies and sources of bias found in the literature that won't be found elsewhere. It is a peer reviewed

summary of the most recent information on polar animals, relative to historical records, based on a review of the 2022 scientific literature and media reports. It is intended for a wide audience, including scientists, teachers, students, decision-makers and the general public.

Executive summary

- There were no reports from either hemisphere in 2022 that would suggest polar wildlife is suffering as a result of reduced sea-ice extent; in both the Arctic and Antarctic, less summer sea ice and increased primary productivity over the last two decades has meant more food for all animals, and explains in part why polar wildlife has been thriving.
- Arctic sea ice in summer has declined since 1979 but the trend flattened after 2007; coverage was again well below average in the Barents and Chukchi Seas in 2022, where continued high primary productivity has provided abundant food resources for wildlife; winter ice coverage in 2022 was slightly lower than 2020, but overall has shown a relatively flat trend since 2011.
- Ice-dependent polar bears worldwide probably now number about 32,000, with a wide range of potential error; a survey of Western Hudson Bay polar bears in 2021 found a population decline of 27% since 2016 that did not correlate with lack of sea ice; a genetically-distinct subpopulation of polar bears was discovered thriving in south-east Greenland; western Barents Sea bears (Norway) are still doing well despite the most profound summer sea ice loss of all Arctic regions.
- Atlantic walrus numbers are still low, but recovering in the Barents Sea and eastern North America; a new population estimate of Pacific walrus in 2019 showed more than 200,000 exist in the Chukchi/Bering Sea area; more killer whales were reported visiting the Eastern Canadian Arctic; in Alaska and the Western Canadian Arctic, bowhead whales are thriving.
- Antarctic sea-ice extent has barely changed since 1979: vital winter ice has slightly increased

overall, while summer ice has slightly declined (with its lowest extent in December 2022), all while overall primary productivity has increased; a new sea-ice predictive model acknowledges previous flaws and does not predict a future decline until 2050 at the earliest.

- Krill are crucial prey for many species (especially huge numbers of great whales and penguins) that live or feed in the Southern Ocean; future intensification of commercial fishing of krill (largely to feed farmed fish) is likely the largest conservation threat to local wildlife, given recent geopolitical tensions over effective fisheries management.
- Numbers of fin, blue, humpback, and southern right whales feeding in Antarctic waters in summer have increased in recent years, and while minke whale numbers appear to have declined, an estimated 500,000 individuals still frequent the region.
- Killer whales (orcas) are the top predator in the Southern Ocean and most populations appear to be thriving; all ice-dependent seals in Antarctica are listed as of 'least concern'.
- Several albatross and large petrel species are considered 'vulnerable' due to deadly interactions with long-line trawlers fishing for Antarctic toothfish (aka Patagonian sea bass), while overfishing of this cod-like species and the herring-like Antarctic silverfish are also a concern.
- Emperor penguins, the largest and most ice-dependent penguin species, were classified as 'Threatened' on the US Endangered Species List in 2022 but remain 'Near Threatened' according to the official 'Red List' because of the large size of their breeding population and uncertainty of future sea-ice predictions, which sea-ice experts acknowledge.



1. Introduction


There were no reports from anywhere around the Arctic or Antarctic in 2022 that would suggest that polar wildlife is suffering as a result of reduced sea-ice extent: no starving polar bears or walrus, no beach-cast dead seals, no marked declines in great whale numbers, no drowned penguin chicks. Inuit in Canada reported more polar bear conflicts with humans, more likely to be an effect of more bears and more people rather than less summer sea ice. The only subpopulation of polar bears that appears to have recently declined did not correlate with any reduction in summer sea ice: some other, unknown, cause was to blame. Contrary to expectations, Antarctic sea ice has been increasing since 1979. Inuit in eastern Canada (Davis Strait) report a marked decline in ringed seal numbers since 1950, but otherwise there were no reports of population declines or reduced health in Arctic seals or walrus. A few Antarctic penguin species and the minke whale appear to have suffered a recent decline in abundance that is unrelated to sea-ice cover. Overall, Antarctic wildlife of all kinds appears to be doing well. In both the Arctic and Antarctic, less summer sea ice and increased primary productivity over the last two decades has meant more food for all animals, and explains in part why polar wildlife has been thriving.

2. The Arctic

Sea ice and primary productivity

According to the NOAA Arctic Report Card 2022, summer sea-ice extent (at September) has declined markedly since 1979 (36.5%), although winter ice levels (at March) have declined very little (9.3%) (Figure 1).⁵ However, in recent years, the trend in March sea-ice coverage has flatlined (since 2011), driven by strong ice production along the Siberian coast; the trend in summer ice extent has been static since 2007.⁶ As a consequence of continued low summer sea-ice extent and reduced ice thickness (which allows beneficial under-ice phytoplankton blooms in summer),

Figure 1: Arctic sea-ice extent, 1979–2022



studies show primary productivity in many regions has continued to climb between 2003 and 2022, especially in the Barents and Chukchi/Bering Seas. Even production of bottom-dwelling phytoplankton has been increasing in some regions.⁷ These phytoplankton blooms provide abundant food for all organisms in the Arctic food chain, including zooplankton ('krill'), benthic invertebrates (such as clams), fish, and marine mammals.⁸

Sea-ice conditions vary regionally across the Arctic in all seasons (Figure 2 shows the locations of the areas discussed). In 2022, winter sea ice at the end of March was low in the Sea of Okhotsk and western Barents Sea but more abundant than usual in Baffin Bay and the Bering Sea. In September, some ice persisted in the Chukchi Sea, which also happened in 2021; however, extent was below normal elsewhere except in the Barents and East Greenland Seas, which had about average ice coverage.⁹



Figure 2: Arctic regions.

Western Hudson Bay (WH) sea-ice coverage and its relationship to polar bears are often considered a 'bellwether' indicator for conditions predicted in a warmer world. Since the abundance of WH polar bears as it relates to length of the ice-free period is now used as a proxy for all other subpopulations, these bears and their sea-ice habitat figure disproportionately in predictions of future species survival and health of the planet in general.¹⁰

In 2022, breakup of sea ice along Western Hudson Bay was about what it had been in the 1980s (11 July $\pm$ 4 days), although a large number of bears tagged a few months earlier by researchers stayed on rapidly-declining sea ice until late August.¹¹ Most bears, even those that came ashore earlier than others, appeared to be in good condition.¹² In the fall, freeze-up was also like it had been in the 1980s (Nov. 16 $\pm$ 5 days).¹³

Summer breakup dates after 2015 have yet to be incorporated into the scientific literature.¹⁴ However, a report by Erin Miller and colleagues in 2022 provided new fall freeze-up dates to 2020: these have been similar to dates in the 1980s (Nov. 16 $\pm$ 5 days) except in 2016 when ice didn't form until the first week of December.¹⁵ The authors found no statistically significant temporal trend in sea ice or departure dates of polar bears from shore between 1991 and 2020, contradicting statements often found in the media that WH fall sea-ice coverage has been steadily declining over the last 30 years.¹⁶

Polar bears


Polar bears are the apex predator of the Arctic. They currently have a relatively large population size, and there has been no change in their historical range since 1979. The International Union for the Conservation of Nature (IUCN), in their 2015 Red List assessment, again listed the polar bear as 'Vulnerable' to extinction, and in 2016, the US upheld its 2008 decision to list the species as 'Threatened' under the US Endangered Species Act.¹⁷ Both decisions were based on computer-modelled future declines, not observed declines. In Canada, where roughly two thirds of entire population lives, polar bears were listed as a species of 'Special Concern' in 2018.¹⁸

The latest IUCN Polar Bear Specialist Group (PBSG) estimate, published in July 2021, is 26,000 (range 22,000–31,000),¹⁹ but took the lowest of several recent estimates available for the Chukchi Sea and ignored figures used in the 2015 IUCN assessment for the Kara and Laptev Seas.²⁰ Survey results postdating the 2015 assessment plausibly bring the mid-point total to about 32,000, with a fairly wide margin of error (Figure 3).²¹ This is much more than the 7,493 (6,660–8,325) bears the public was assured would be all that would remain²² given the sea-ice conditions since 2007.²³

For a detailed discussion of the status of all 19 subpopulations, see the *State of the Polar Bear Report 2020*.²⁴ In 2022, the official government report on the 2017–2018 Davis Strait subpopulation survey was finally made available.²⁵ It confirmed

Figure 3: Estimates of the global polar bear population

1960 to date. The 1981, 1993, and 2015 estimates are from the IUCN PBSG, 1960 from Crockford 2019 (pp. 102–105) and Anonymous 1966 (p. 11), and the 2021 estimate is from Crockford 2022.



earlier news reports of an estimate of 2,015 bears (range 1,603–2,588), statistically indistinguishable from the 2007 estimate of 2,158 (range 1,833–2,542). What hadn't been made public by the media earlier was that no correlation had been found between polar bear survival and sea-ice conditions (e.g. neither length of ice-free period or mean summer sea-ice concentration). An increased harvest rate *was* considered to be a concern. Other evidence – especially bears in better condition in recent years compared to 2007 – suggest the Davis Strait population may be stable or even increasing, although the authors suggest a stated Nunavut management objective to reduce the subpopulation had been successful.²⁶

Also in 2022, some results of an aerial survey of Western Hudson Bay in late summer 2021 were released to the media, but the government report with full details was withheld. Although a 27% decline in numbers since 2016 was reported (618, compared to 842 in 2016), it was emphasised that this phenomenon did not correlate with a decline in WH sea-ice coverage: instead, an unexplained lack of ringed seal prey was suggested as an explanation.²⁷

Overall, bears in virtually all regions have been reported to be in good condition in recent years. Improved body condition has been reported for Davis Strait bears in 2017/2018 (compared to those examined a decade before) and more recently for bears in M'Clintock Channel, Gulf of Boothia, and Kane Basin in the Canadian Arctic, confirmed by on-the-ground experiences of Inuit in Davis Strait and Gulf of Boothia.²⁸ In the Svalbard region of the Barents Sea, the body condition of male bears in the spring of 2022 was below what it had been in 2021 and well below 2019, but well within the natural range of variation since 1993 (conditions in 2003 were worse).²⁹

Despite concerns that continued low summer sea-ice coverage would precipitate more conflicts between polar bears and people, it hasn't quite worked out as expected.³⁰ While Inuit in Canada, especially, have noted more problems with bears compared with previous decades, they usually attribute this phenomenon to more bears, not less ice, and some residents have pointed out that polar bears tend to be more aggressive when their population densities are high.³¹ A perceived increase in problems with bears visiting garbage dumps across the Arctic in recent years has also been blamed on hunger, although with little evidence.³² An alternative explanation is an increase in human population (and thus more garbage) and higher polar bear numbers over the last six decades (Figure 3).³³

Overall, recent data collected from across the Arctic, but especially in the western Barents Sea and Western Hudson Bay, do not support the assumption, stated repeatedly by scientists in the field, that summer sea-ice loss inevitably leads to reduced polar bear survival: in WH, a population decline of 27% over five years was not associated with any lack of ice, and in Svalbard, polar bears are still doing well despite the greatest summer sea-ice loss of all Arctic regions.³⁴

Walrus, seals, and whales

Ringed seals (*Pusa hispida*) and bearded seals (*Erignathus barbatus*), and particularly their pups, are critical prey of polar bears across the Arctic.³⁵ In some regions, other seal species, walrus (*Odobenus rosmarus*), beluga (*Delphinapterus leucas*), and narwhal (*Monodon monoceros*) are consumed as well,³⁶ and bears may also scavenge the carcasses of bowhead whales (*Balaena mysticetus*) that have died naturally or as a consequence of indigenous hunting.³⁷ These species are also important food resources of indigenous Arctic peoples, and healthy populations are critical to human survival in the Arctic.

Walrus


In 2017, the US Fish and Wildlife Service determined that the Pacific subspecies of walrus (*O. r. divergens*) no longer required the statutory safeguards it had enjoyed since the Marine Mammal Protection Act of 1972.³⁸ In 2022, the latest population assessment for Pacific walrus in the US portion of the Chukchi Sea was published, with the total put at 189,000 (135,000–251,000) in 2019, based on animals counted at Point Lay, Alaska in late summer.³⁹ As this 'most accurate to date' estimate did not include animals located in the Russian portion of the Chukchi Sea or the southern Bering Sea at the time of the survey,⁴⁰ it seems likely the entire Pacific walrus population count is well over 200,000, despite continued summer sea-ice loss in the region. This is similar to an estimate from 2015 (283,213; range 93,000–478,975).⁴¹

In contrast, the Atlantic walrus subspecies (*O. r. rosmarus*) has not rebounded from historical overhunting to nearly the same extent, but the numbers are recovering regardless: it was

classified by the IUCN in 2016 as 'near threatened', an important step below 'vulnerable', which took into account healthy population numbers (about 25,000 at that time). Even as summer sea ice declined markedly, the Svalbard population increased by 109% between 2006 and 2018 (and 42% between 2012 and 2018), for a final estimate of 5,503 (Figure 4).⁴² However, since walrus are notoriously difficult to count accurately, this is almost certainly an underestimate.⁴³ In neighbouring Russian territory of Franz Josef Land, walrus numbers in 2017 were estimated to be approaching 'pre-hunting levels' of 9,000–11,000.

Figure 4: Walrus population in Svalbard since 1980

The numbers on the graph start with two baseline estimated numbers provided in 1980 and 1993 (1980 = 100 and 1993 = 741), and conclude with population counts from aerial surveys done in 2006, 2012, and 2018 (2006 = 2629; 2012 = 3886; 2018 = 5503). From Norwegian Polar Institute 2022d.



On the other side of the Atlantic, Inuit in Eastern Canada have noticed that walrus seem to be moving seasonally in response to sea-ice changes, but consider the walrus to be 'not at risk', given an estimated 21,400 individuals in 2021. In spite of this, the Canadian government Species at Risk Act listed the Atlantic walrus as of 'special concern' in 2017.⁴⁴

Seals

A report by Nunavut Inuit in the Davis Strait region of eastern Canada revealed that (Figure 5):


- Numbers of bearded seals have been unchanged over the last seven decades.
- Harp seal and polar bear numbers have steadily increased from very low abundance.
- Ringed seals, which used to be very abundant, have steadily declined.⁴⁵

While a marked decline in ringed seal numbers in Davis Strait since the 1950s has not been reported by biologists, it appears that no studies have been done. Furthermore, as this region connects via Hudson Strait to critical polar bear habitat in Hudson Bay, a region-wide decline in ringed seal numbers might explain

Figure 5: Polar bear and seal encounters 1940–2019

Inuit perceptions of changes in animal populations in Davis Strait (eastern Canada), based on recollections of encounters in the locale. Suggested as a proxy for relative species abundance in the area.

- Bearded seal
- Ringed seal
- Harp seal
- Polar bear



the recent polar bear population decline in Western Hudson Bay, as discussed above. Possibilities offered by Inuit to explain the decline in ringed seals included: an increase in polar bear predation (especially during the pupping season); an increase in community harvests; ringed seals moving to other areas (either following prey, or due to displacement by other species); lack of sea ice; warmer ocean waters; increase in fox predation; and changing winds.⁴⁶

Whales

Although some scientists are worried about the future of narwhal and beluga in the Arctic, at this time there is no evidence that most subpopulations of these species are struggling due to reduced sea-ice coverage in summer.⁴⁷ On the other hand, research on bowhead whales in the western Arctic has revealed a marked increase in numbers over the past 30 years, despite recent sea-ice changes and aboriginal subsistence hunting. The study authors suggest this is likely due to increases in ocean primary productivity.⁴⁸

Historically, killer whales (orcas, *Orcinus orca*) have been rare visitors to the Arctic. However, scientists have noted an increase in sightings, especially in the Cumberland/Lancaster Sound region of the eastern Canadian Arctic and in Hudson Bay, which seems to be correlated with reduced summer sea ice. However, the same phenomenon has been noted in the Gulf of St Lawrence (which is always ice-free in summer), indicating that the influx of the predatory whales might have another cause (Figure 5).⁴⁹ Perhaps not coincidentally, in the summer, thousands of pregnant beluga whales* (known to be preyed upon by killer whales) use the Cumberland/Lancaster Sound area in the High Arctic and western Hudson Bay, especially to give birth and nurse their newborns, while a few hundred reside year-round in the

* Note that belugas inhabit Arctic and sub-Arctic waters of North America and Eurasia but oddly not Eastern Greenland.

Figure 6: Summary of killer whale population movements

1. Lancaster Sound, Cumberland Sound. 2. Hudson Bay. 3. St Lawrence Estuary. Reference: Halliday et al. 2022.



St. Lawrence River estuary.⁵⁰ Contrast this relative rarity of killer whale sightings in the Arctic during the summer to the tens of thousands estimated to frequent the Southern Ocean, including pack ice habitat *in winter* (Section 7). This suggests these predatory whales do not have an inherent aversion to sea ice.

Birds and fish

Birds

According to a new report in 2022, most geese species that migrate into the Arctic to breed and raise their chicks remain abundant, with either increasing or stable trends.⁵¹ Lesser snow geese (*Anser caerulescens caerulescens*) and Aleutian Canada geese (*Branta hutchinsii*, a small, formerly endangered subspecies) in the Western Arctic (Bering/Chukchi Seas) have been doing especially well in recent years.⁵² A report that 450 sea birds died in the Bering and southern Chukchi Seas this year was well below previous years.⁵³ The authors admit they don't really know the cause of these die-offs but suggest it is most likely to be starvation. A US Fish and Wildlife Service report on the die-off phenomenon in 2016 noted that sea birds can starve to death if they haven't eaten in as few as four days, which makes them vulnerable to any disruption of feeding, especially during storms.⁵⁴

Fish

Some species of bottom-dwelling invertebrates and fish have been documented expanding north into the Arctic from their usual locations in temperate waters (notably in the Barents, Bering, and Beaufort Seas), but it is clear that such changes cannot be blamed exclusively on warming caused by human fossil fuel use and may not be permanent.⁵⁵ So-called 'regime shift' changes have occurred naturally in the past: sea-ice changes along the coast of East Greenland are well documented, as are changes in eastern Canada, the Bering Sea, and the Svalbard region of the Barents Sea.⁵⁶

3. The Antarctic

Sea ice and primary productivity

In contrast to the Arctic, which is an ocean basin largely surrounded by land, the Antarctic is a frigid continent surrounded by an ocean that freezes extensively in winter. However, usually, very little sea ice persists over the austral summer because it is virtually all thin, first year ice (around 1 m thick at most).⁵⁷ Antarctic winter sea-ice area (at September) has been increasing slowly since 1978 at a rate of about 1.7% per decade, especially in the Ross Sea,⁵⁸ and summer ice levels (at February) have declined slightly (Figure 7).

On 1 September 2021, winter sea ice reached its annual maximum. This was both a record high (18.75 mkm²) and the second-earliest annual peak on record, likely due to the second-lowest temperatures on record for June-July-August. In 2022, maximum winter ice extent was back down to 18.19 mkm² (at 16 September).⁵⁹ According to sea-ice experts, these latest readings exemplify the extreme inter-annual variability that has been documented over the last decade, both for overall ice extent and regional extent around the continent. The greatest maximum extent recorded since 1978 was in 2014 (20.11 mkm² at 22 September; *average* for that month was 19.76 mkm²), although 2013 and 2012 were almost as high (Figure 7).⁶⁰ A recent paper has found that along the Antarctic Peninsula, '85% of the ice shelf perimeter...has advanced since the early 2000s, in contrast to the extensive retreat of the previous two decades,' which they attribute in part to a stabilising effect from increased sea-ice extent.⁶¹

Stable Antarctic ice extent over the last few decades contradicts the expectations of global sea-ice loss predicted by climate models.⁶² Moreover, a new model published in 2022 does not predict significant winter sea-ice loss in the Southern Ocean until about 2050 – more than three additional decades of stable sea-

Figure 7: Antarctic sea ice extent, 1979 to date.

Monthly averages for September (max) and February (min). From Colucci 2022 <https://www.severe-weather.eu/global-weather/antarctic-sea-ice-extent-record-low-anomaly-observed-rrc/>.




Figure 8: Typical summer sea ice extent in the Southern Ocean

Shown here for February 2020



ice cover.⁶³ Unfortunately, such long-overdue revisions to sea-ice projections have not been incorporated into the biological models used to predict survival and conservation status of Antarctic species dependent on sea ice. Their populations are thus still predicted to decline over time.⁶⁴

Satellite observations indicate continued strong primary productivity in most regions of the Southern Ocean over the last two decades, especially in the Ross Sea and off the coast of the Antarctic Peninsula, but also in coastal polynyas (areas of open water surrounded by ice) around the entire continent.⁶⁵ As in the Arctic, phytoplankton blooms provide abundant food for all organisms in the food chain, but especially krill (*Euphausia superba*), the shrimp-like crustacean that is the foundation of the Antarctic ecosystem.⁶⁶ There are several hot-spots of krill abundance: a huge one in the Weddell Sea and several others scattered around the continent.

Krill are preyed upon by hundreds of thousands of whales that converge on the Southern Ocean every summer. Several important fish, penguin, and seal species also depend on them, as do squid and flying sea birds. Humans now do as well. Trawlers from China, Norway, South Korea, Ukraine and Chile fish for krill commercially, and while their catch has long been restricted to a maximum of 620,000 tons annually – about 1.6% of the estimated 400 million tons (range 300–500t) total biomass – they have never yet removed that much (e.g. the largest catch of 450,000 tons was reached in 2020).⁶⁷

Whales and seals

Whales

In the Antarctic, the top of the food chain is occupied by the

killer whale. Killer whales hunt like packs of wolves, consuming small whales, seals, penguins, and fish. Some prowl the Southern Ocean at the edge of the sea ice in summer (where many of their prey species congregate to feed) but others penetrate the pack ice to hunt year round. Four ecologically and genetically distinct 'ecotypes' of these killer whales exist, each with a distinctive overall size and variant of the species' typical facial markings (e.g. eye patch).⁶⁸

- Type A, the largest (males ~8.0–9.5 m), hunts in the ice-free waters of the Southern Ocean just beyond the pack ice during the austral summer, feeding primarily on diminutive Antarctic minke whales (*Balaenoptera bonaerensis*) and calves of larger whale species.
- Type B1, aka the 'pack ice killer whale,' is smaller (males ca. 7.8 m) and eats primarily Weddell seals (*Leptonychotes weddellii*), which groups cooperatively to 'wave-wash' prey off small ice floes.
- Type B2, the Gerlache killer whale, is smaller still (males ~6.4 m) and often hunts in large groups in open water off the Antarctic Peninsula in summer, primarily for small penguins (especially gentoos and chinstraps), but perhaps also fish.
- Type C, the 'dwarf killer whale,' is the smallest (males 6.0 m, females 5.5 m); it hunts deep in the pack ice and prowls cracks in the landfast ice of the Ross Sea, feeding primarily on large Antarctic toothfish (Chilean sea bass, *Dissostichus mawsoni*).

Types B2 and C appear to be year-round residents, as both have been observed in the pack ice in winter and summer.⁶⁹ Killer whale type B2 (the small, penguin-eater) seems to be overall about twice as abundant as types A and type B1, although around the Antarctic Peninsula in 2017/2018 it was found to be more than seven times as common (~102 for B1 – apparently a slight decline since 2010 – vs ~740 for B2, statistically similar to estimates in 2010).⁷⁰ Over a similar period (ending 2016/2017), type A killer whales increased significantly in abundance (from ~91 to ~149).⁷¹

The Southern Ocean has the largest concentration of orcas worldwide, and killer whales are the second-most abundant whale found off Antarctica, after minke whales.⁷² A series of surveys in 2001 estimated the total killer whale population in Antarctic waters in the summer was ~25,000–27,000, but this estimate was likely an underestimate at the time given that survey vessels did not venture into the pack ice that many orcas frequent.⁷³ In 2020, the Australian Antarctic Program estimated their abundance at 70,000.⁷⁴

Perhaps it's not a coincidence (given the growing abundance of predatory Type A killer whales) that numbers of Antarctic minke whales have apparently declined in recent years. The species currently numbers about 500,000, down from about 720,000 in earlier assessments (an apparent 31% decline). Although there

are concerns about the accuracy of these figures, in part due to the tendency of this species to enter the pack ice (where survey ships seldom enter), the species was listed as 'near threatened' in 2018 by the IUCN.⁷⁵

Other species of Antarctic whales are recovering from historic industrial hunting of whales, many of which congregate off the extremely productive Antarctic Peninsula to feed in the austral summer. Southern fin whales (*Balaenoptera physalus quoyi*) numbered about 8,000 individuals in 2019 and have recently established large feeding aggregations of 50–150 individuals, which they did before historic whaling decimated their populations; globally, the species is still listed as 'vulnerable' by the IUCN, even though numbers are increasing, because the population is much smaller than it was before modern whaling.⁷⁶ The International Whaling Commission (IWC) determined that the population size of humpback whales (*Megaptera novaeangliae*) in the Southern Hemisphere had rebounded to about 70% of pre-hunting levels.⁷⁷

There are estimated to be approximately 3–4,000 southern right whales (*Eubalaena australis*) in the southern hemisphere, a slow recovery from near extinction due to over-hunting prior to 1935. It is classified as of 'least concern' by the IUCN.⁷⁸ In contrast, Antarctic blue whales (*Balaenoptera musculus intermedia*), which often enter the pack ice, are also slowly recovering (estimated 6,500 individuals, range 5,000–8,000), but were considered 'critically endangered' by the IUCN in 2018 because their numbers comprised only about 2.5% of their pre-exploitation abundance.⁷⁹

Seals

There are only four species of ice-dependent seals in the Antarctic, versus six in the Arctic (seven if you count the walrus). Southern elephant seals (*Mirounga leonina*) and Antarctic fur seals (*Arctocephalus gazella*) are not ice-dependent, although they do forage in Antarctic offshore productive zones, including the pack ice, even in winter.⁸⁰

Weddell seals are among the largest seals (~2.5–3.5 m long; weight 400–600 kg) and have been observed diving to 600 m. They primarily eat squid and fish, especially Antarctic toothfish and silverfish (*Pleuragramma antarcticum*). Weddell seals make their breeding and pupping colonies on stable landfast ice in the late winter/spring (September–November) but move into the pack ice during the austral summer to feed. Uniquely among seals, females may occasionally produce twins. Weddell seals are the second most abundant species of Antarctic phocid, after the crabeater seal: the total population was estimated at about 300,000 individuals in 2011.⁸¹ The IUCN classified the crabeater as being of 'least concern' in 2015, due to 'widespread occurrence, large population size and lack of major threats'.⁸²

Leopard seals (*Hydrurga leptonyx*), Ross seals (*Ommatophoca rossii*) and crabeater seals (*Lobodon carcinophagus*) all breed and

pup within the mobile pack ice and all are listed as of 'least concern' by the IUCN. Crabeaters are the most abundant seal in the Southern Ocean, with estimates of at least 7 million animals.⁸³ They do not in fact eat crabs, but are voracious consumers of krill, which they strain out of the water with specialized multi-lobed teeth. Their primary predators are leopard seals, which take a large percentage of young crabeaters each year. Leopard seals are about the same size as Weddell seals, and abundance is estimated at 220,000–440,000 animals. Young leopard seals primarily eat krill but adults switch to eating young seals, fur seals, and penguins.⁸⁴ Ross seals, which eat squid and fish, are the least common species and their abundance is uncertain.⁸⁵

Penguins

Several species of penguins breed on sub-Antarctic islands but feed in Antarctic waters during the non-breeding season, including macaroni (*Eudyptes chrysolophus*), southern rockhopper (*Eudyptes chrysocome*) and king (*Aptenodytes patagonicus*). Of these the macaroni penguin is the most abundant (estimated at 13 million adults), but is declining (IUCN, 'vulnerable'), southern rockhoppers (estimated at 2.5 million adults) have recently declined due to habitat loss, human disturbance and competition from commercial fishing (IUCN, 'vulnerable'), while king penguin numbers are stable at about 2.2 million adults (IUCN, 'least concern').⁸⁶

Gentoo (*Pygoscelis papua*), chinstrap (*Pygoscelis antarctica*), and Adélie (*Pygoscelis adeliae*) penguins require ice-free ground for nesting in summer: gentoos and chinstraps choose sub-Antarctic habitats for nesting, but feed in Antarctic waters year-round, while Adélies prefer ice-free ground along the Antarctic coast in summer and pack ice in winter.⁸⁷ Gentoo populations are considered stable (IUCN, 'least concern'), with a population of about 774,000 adults.⁸⁸ Adélies are currently increasing in abundance (IUCN, 'least concern'), with population estimates of 14–16 million individuals.⁸⁹ Chinstrap numbers have declined in some regions but increased in others (overall abundance, about 8 million adults) and while it is listed as 'least concern' by the IUCN, it is considered at risk for possible overall declines.⁹⁰ There are concerns that chinstrap penguins in particular are vulnerable to future declines in krill abundance anticipated to occur over the next century due to predicted sea-ice loss due to human-caused climate change (but see Section 6 on sea ice).⁹¹

Emperor penguins (*Aptenodytes forsteri*, up to 100 cm tall) are the only species that breeds, lays eggs, and feeds its chicks to fledgling stage on landfast ice in Antarctica (Figure 8). Also unique to this species, it conducts these physiologically demanding tasks over the frigid winter and spring months (Figure 9). Consequently, emperor penguins are the only species apt to be directly affected by annual or decadal variations in Antarctic sea-ice extent.

Emperor penguin populations in 2019 were found to have grown by up to 10% since 2009 – to as many as 282,150 breed-

Figure 9: Breeding range of the emperor penguin

A few sites are well offshore. Red circles are the new sites reported in 2020 by Fretwell and Trathan. Major ice shelves are named.





Figure 10: Emperor penguin breeding cycle.

From Amos, BBC 5 Aug 2020.



ing pairs (up from about 256,500) out of a total population of over 600,000 birds – despite a loss of thousands of chicks in 2016 when an ice shelf collapsed in the Weddell Sea.⁹² Nevertheless, biologists studying this species petitioned the IUCN to upgrade emperor penguins from ‘near threatened’ (at 2018) to ‘vulnerable’⁹³, based on models that use the most extreme (and therefore highly unlikely) RCP8.5 ‘worst case’ climate change scenario.⁹⁴ Not only do these extreme models implausibly suggest emperor penguins could be close to extinction by 2100, they also fail to account for the fact that recent sea-ice loss has not been nearly as severe as expected, further calling into doubt the accuracy of these predictions.⁹⁵

In 2019 the IUCN declined to change its 2018 assessment, stating: ⁹⁶

This species is listed as near threatened because it is projected to undergo a moderately rapid population decline over the next three generations owing to the projected effects of climate change. However, it should be noted that there is considerable uncertainty over future climatic changes and how these will impact the species.

So far, the localised die-off of chicks in one of the many colonies along the edge of the Weddell Sea in 2016 has not recurred. However, in 2022 the US Fish and Wildlife Service responded to pressure and formally listed the emperor penguin as ‘threatened’ under the Endangered Species Act.⁹⁷

Other birds and fish

Birds

A variety of albatrosses, petrels, skuas, fulmars, shearwaters, gulls, and terns, nest in summer on sub-Antarctic islands or at ice-free locations around the Antarctic continent (especially the Antarctic Peninsula) and feed in the rich offshore waters of the Southern Ocean. Several albatross and large petrel species are considered ‘vulnerable’ by the IUCN due to their penchant for stealing baited hooks set by long-line trawlers fishing for Antarctic toothfish.⁹⁸

Fish

The Antarctic toothfish is a cod-like species that can reach lengths of over 2 metres and, because it is fished commercially, has generated concerns of over-harvesting.⁹⁹ The toothfish eats krill and Antarctic silverfish, and is preyed upon by some killer whales and Weddell seals. The small schooling Antarctic silverfish (aka ‘Antarctic herring’) is also an important prey species for some killer whales, humpback whales, minke whales, Weddell seals, and at least two species of penguins (Adélies and emperors).¹⁰⁰

4. Discussion

Contrary to expectations, while Arctic sea-ice extent has seen an overall decline since 1979, that trend has stalled in recent years: since 2007 for summer ice and about 2011 for winter ice. Also

contrary to expectations, the open-water season in Hudson Bay has not continued to increase since a 'step change' took place in 1998. This explains why a recent decline in polar bear abundance in Western Hudson Bay did not correlate with lack of sea ice. However, summer ice has declined over the Barents Sea to a much greater extent than any other Arctic region since at least 2003, and this has had the opposite effect expected for seal and polar bear populations because of the increased primary productivity that a longer open-water season promotes: Barents Sea polar bears are thriving. A similar phenomenon has been documented in the Chukchi Sea: less summer ice has meant more seals and abundant, healthy bears.

In contrast, Antarctic sea-ice area has been surprisingly stable: overall winter ice cover has increased slightly since 1979. Perversely, climate models used to forecast future conditions (assumed to be driven by ever-increasing human-caused carbon dioxide emissions) predicted marked declines in Antarctic sea ice over the recent decades and even more over the 21st century.

Sea-ice experts have documented their concerns about this discrepancy between observations and model predictions for more than ten years; in 2022 a model proposed that virtually stable Antarctic winter sea-ice cover could be expected to continue to at least 2050, with only modest declines after that. However, this evidence has apparently been ignored by biologists proclaiming future declines in abundance of Antarctic species (especially krill, and several penguin species).

This is especially apparent for the emperor penguin, for which the most pessimistic models predict near-extinction by 2100 due to winter sea-ice loss: not only are the 'worst-case' scenarios that underlie these claims implausible (due to far-fetched assumptions of future carbon dioxide emissions), but the sea-ice factor is now also known to be quite flawed. It is apparent that emperor penguins were listed in the US (in late 2022) as 'threatened' based on predicted winter sea-ice loss using a justification that ignored well-founded evidence – published in the scientific literature by reputable sea-ice researchers – that the sea-ice models used were not fit for purpose.

The same flawed sea-ice models underlie claims of marked future declines in Antarctic krill abundance, and of resulting species collapse across the continent, a function of the central role that krill plays in Antarctic ecology. All such predictions should therefore be considered flawed.

A decline in winter sea ice blamed on climate change cannot be the primary driver of changes in Antarctic faunal abundance, simply because winter ice extent has not declined. Chinstrap penguins and minke whales, for example, appear to have suffered recent declines in abundance but

those changes are unrelated to a loss of winter sea-ice cover.

Overall, Arctic and Antarctic wildlife of all kinds appeared to be doing well, with very few exceptions. Polar bear, Atlantic walrus, and bowhead numbers in the Arctic continue to recover from overhunting, as do fin, humpback, blue, and southern right whale numbers in the Antarctic (although some more slowly than others).

There were no reports from either hemisphere in 2022 that would suggest polar wildlife is suffering as a result of reduced sea-ice extent (in summer for the Arctic, winter for the Antarctic): no news of starving polar bears or walrus, beach-cast dead seals, or drowned penguin chicks. In both the Arctic and Antarctic, less summer sea ice and increased primary productivity over the last two decades has meant more food for all animals, which explains why polar wildlife has been thriving.

Bibliography

- Aars, J., Lunn, N. J. and Derocher, A.E. 2006. Polar Bears: Proceedings of the 14th Working Meeting of the IUCN/SSC Polar Bear Specialist Group, Seattle, Washington, 20–24 June 2005. Occasional Paper of the IUCN Species Survival Commission 32. IUCN, Gland, Switzerland.
- Aars, J., Marques, T.A., Lone, K., et al. 2017. The number and distribution of polar bears in the western Barents Sea. *Polar Research* 36:1, 1374125.
- Agence-France Presse. 2022. 'Polar bears vanishing from "polar bear capital of the world" in Canada. *The Guardian* (UK), 24 December. <https://www.theguardian.com/world/2022/dec/24/polar-bears-vanishing-from-polar-bear-capital-of-the-world-in-canada>.
- Amos, J. 2019. 'Climate change: Emperor penguin 'needs greater protection'. BBC News, 9 October. <https://www.bbc.com/news/science-environment-49978007>.
- Amos, J. 2020. 'Climate change: Satellites find new colonies of Emperor penguins.' BBC News (UK), 5 August. <https://www.bbc.com/news/science-environment-53549299>.
- Amstrup, S.C. 2003. Polar bear (*Ursus maritimus*). In: *Wild Mammals of North America*, G.A. Feldhamer, B.C. Thompson and J.A. Chapman (eds), pg. 587–610. Johns Hopkins University Press, Baltimore.
- Amstrup, S.C., Marcot, B.G. and Douglas, D.C. 2007. Forecasting the rangewide status of polar bears at selected times in the 21st century. Administrative Report, US Geological Survey. Reston, Virginia.
- Andersen, M., Kovacs, K.M. and Lydersen, C. 2021. Stable ringed seal (*Pusa hispida*) demography despite significant habitat change in Svalbard, Norway. *Polar Research* 40:5391. <http://dx.doi.org/10.33265/polar.v40.5391>.
- Arce, F., Hindell, M.A., McMahon, C.R., et al. 2022. Elephant seal foraging success is enhanced in Antarctic coastal polynyas. *Proceedings of the Royal Society, Biological Sciences B* 289(1967):20212452.
- Ardyna, M. Mundy, C.J., Mayot, N., et al. 2020a. Under-ice phytoplankton blooms: shedding light on the "invisible" part of Arctic primary production. *Frontiers in Marine Science* 7:608031. <https://doi.org/10.3389/fmars.2020.608032>.
- Ardyna, M. Mundy, C.J., Mills, M.M., et al. 2020b. Environmental drivers of under-ice phytoplankton bloom dynamics in the Arctic Ocean. *Elementa Science of the Anthropocene* 8:30. <https://doi.org/10.1525/elementa.430>.
- Atkinson, S.N., Laidre, K.L., Arnold, T.W., et al. 2021. A novel mark-recapture-recovery survey using genetic sampling for polar bears *Ursus maritimus* in Baffin Bay. *Endangered Species Research* 46:105–120.
- Australian Antarctic Program 2020. 'Killer whale.' Australian Antarctic Program, 31 August. <https://www.antarctica.gov.au/about-antarctica/animals/whales/killer-whale/>.
- Australian Antarctic Program 2022. 'Blue whale.' Australian Antarctic Program, 6 September. <https://www.antarctica.gov.au/about-antarctica/animals/whales/blue-whale/>.
- Bailleau, F., Charrassin, J.-B., Ezraty, R. et al. 2007. Southern elephant seals from Kerguelen Islands confronted by Antarctic sea ice: changes in movements and diving behaviour. *Deep Sea Research Part II: Topical Studies in Oceanography* 54(3–4):343–355.
- Beatty, W.S., Lemons, P.R., Sethi, S.A., et al. 2019. Estimating Pacific walrus abundance and demographic rates from genetic mark-recapture. US Department of the Interior, Bureau of Ocean Energy Management, Anchorage, Alaska. OCS Study BOEM 2019-059.
- Bengtsson, J.L. 2009. Crabeater seal: *Lobodon carcinophaga*. *Encyclopedia of Marine Mammals* (2nd edition):290–292.

- Bengtsson, O., Lydersen, C., Kovacs, K.M., et al. 2020. Ringed seal (*Pusa hispida*) diet on the west coast of Spitsbergen, Svalbard, Norway: during a time of ecosystem change. *Polar Biology* 43:773–788. <https://doi.org/10.1007/s00300-020-02684-5>.
- Bengtsson, O., Hamilton, C.D., Lydersen, C., et al. 2021. Distribution and habitat characteristics of pinnipeds and polar bears in the Svalbard Archipelago, 2005–2018. *Polar Research* 40:5326. <http://dx.doi.org/10.33265/polar.v40.5326>.
- Benson, A.J. and Trites, A.W. 2002. Ecological effects of regime shifts in the Bering Sea and eastern North Pacific. *Fish and Fisheries* 3:95–113.
- Bestley, S., Andrews-Goff, V., van Wijk, E., et al. 2019. New insights into prime Southern Ocean forage grounds for thriving Western Australian humpback whales. *Scientific Reports* 9: 13988.
- Bilby, E. 2022. 'Cloistered Arctic whales face a bigger climate threat than polar bears'. *European Commission*, 29 December. <https://ec.europa.eu/research-and-innovation/en/horizon-magazine/cloistered-arctic-whales-face-bigger-climate-threat-polar-bears>.
- BirdLife International. 2018. *Aptenodytes forsteri*. The IUCN Red List of Threatened Species 2018: e.T22697752A132600320. Downloaded on 07 August 2020. <https://www.iucnredlist.org/species/22697752/132600320>.
- BirdLife International. 2020a. *Aptenodytes forsteri*. The IUCN Red List of Threatened Species 2019: e.T22697752A132600320. Downloaded on 26 October 2022. <https://www.iucnredlist.org/species/22697752/157658053>.
- BirdLife International. 2020b. *Pygoscelis adeliae*. The IUCN Red List of Threatened Species 2020: e.T22697758A157660553. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22697758A157660553.en>.
- BirdLife International. 2020c. *Pygoscelis antarcticus*. The IUCN Red List of Threatened Species 2020: e.T22697761A184807209. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22697761A184807209.en>.
- BirdLife International. 2020d. *Pygoscelis papua*. The IUCN Red List of Threatened Species 2020: e.T22697755A157664581. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22697755A157664581.en>.
- BirdLife International. 2020e. *Eudyptes chrysocome*. The IUCN Red List of Threatened Species 2020: e.T22735250A182762377. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22735250A182762377.en>.
- BirdLife International. 2020f. *Eudyptes chrysolophus*. The IUCN Red List of Threatened Species 2020: e.T22697793A184720991. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22697793A184720991.en>.
- BirdLife International. 2020g. *Aptenodytes patagonicus*. The IUCN Red List of Threatened Species 2020: e.T22697748A184637776. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T22697748A184637776.en>.
- Blanchard-Wrigglesworth, E., Eisenman, I., Zhang S. et al. 2022. New perspectives on the enigma of expanding Antarctic sea ice. *Eos* 103. <https://doi.org/10.1029/2022EO220076>.
- Blanchard-Wrigglesworth, E., Roach, L.A., Donohoe, A. and Ding, Q. 2021. Impact of winds and Southern Ocean SSTs on Antarctic sea ice trends and variability. *Journal of Climate* 34(3):949–965.
- Bornemann, H., Kreyscher, M., Ramdohr, S. et al. 2000. Southern elephant seal movements and Antarctic sea ice. *Antarctic Science* 12(1):3–15.
- Bromaghin, J.F., Douglas, D.C., Durner, G.M., et al. 2021. Survival and abundance of polar bears in Alaska's Beaufort Sea, 2010–2016. *Ecology and Evolution* 11(20):14250–14267. <https://doi.org/10.1002/ece3.8139>.
- Brooks, C.M., Ainley, D.G., Jacquet, J. et al. 2022. Protect global values of the Southern Ocean ecosystem, *Science* 378(6619):477–479.

- Brown, T.A., Galicia, M.P., Thiemann, G.W., et al. 2018. High contributions of sea ice derived carbon in polar bear (*Ursus maritimus*) tissue. *PLoS One* 13(1):e0191631. <https://doi.org/10.1371/journal.pone.0191631>.
- Calvert, W. and Stirling, I. 1990. Interactions between polar bears and overwintering walruses in the Central Canadian High Arctic. *Bears: Their Biology and Management* 8:351–356.
- Campana, S.E., Stefánsdóttir, R.B., Jakobsdóttir, K. et al. 2020. Shifting fish distributions in warming sub-Arctic oceans. *Scientific Reports* 10:16448.
- Carneiro, A.P.B., Clark, B.L., Pearmain, E.J., et al. 2022. Fine-scale associations between wandering albatrosses and fisheries in the southwest Atlantic Ocean, *Biological Conservation* 276:109796.
- Castro de la Guardia, L., Myers, P.G., Derocher, A.E., Lunn, N.J., Terwisscha van Scheltinga, A.D. 2017. Sea ice cycle in western Hudson Bay, Canada, from a polar bear perspective. *Marine Ecology Progress Series* 564: 225–233.
- CBC News 2021. 'Orca whales splash cod fishing excursion near St. Philip's.' *CBC News*, 5 July. <https://www.cbc.ca/news/canada/newfoundland-labrador/nl-orca-whales-cod-fishing-1.6091056>.
- Chakilev, M.V. and Kochnev, A.A. 2019. Monitoring results of the Pacific walrus (*Odobenus rosmarus divergens*) haulout site at Cape Serdtse-Kamen (Chukchi Sea) in 2016–2017. In, *Marine Mammals of the Holarctic, Papers of the Tenth International Conference (29 October-2 November 2018)*, Volume 1, pg. 381–391. Marine Mammal Council, Moscow. [Russian and English].
- Chinchar, A. 2021. 'Antarctica's last 6 months were the coldest on record.' *CNN*, 9 October. <https://www.cnn.com/2021/10/09/weather/weather-record-cold-antarctica-climate-change/index.html>.
- Christie, F.D.W., Benham, T.J., Batchelor, C.L., et al. 2022. Antarctic ice-shelf advance driven by anomalous atmospheric and sea-ice circulation. *Nature Geoscience* 15:356–362.
- Chua, M., Ho, S.Y.W., McMahon, C.R. et al. 2022. Movements of southern elephant seals (*Mirounga leonina*) from Davis Base, Antarctica: combining population genetics and tracking data. *Polar Biology* 45:1163–1174.
- Collins, M.A., Hollyman, P.R., Clark, J. et al. 2021. Mitigating the impact of longline fisheries on seabirds: lessons learned from the South Georgia Patagonian toothfish fishery (CCAMLR Subarea 48.3). *Marine Policy* 131:104618.
- Collins, S. 2022. 'Sea ice can control Antarctic ice sheet stability, new research finds.' *University of Cambridge press release*, 13 May. <https://www.cam.ac.uk/stories/sea-ice-controls-ice-sheet-stability>.
- Colucci, R.R. 2022. 'Antarctic Sea ice extent hits a new record low, possibly heading to set the most significant negative anomaly ever observed in the satellite era.' *Severe Weather EU*, 31 December. <https://www.severe-weather.eu/global-weather/antarctic-sea-ice-extent-record-low-anomaly-observed-rrc/>.
- Comiso, J.C., Gersten, R.S., Stock, L.V., et al. 2017. Positive trend in the Antarctic sea ice cover and associated changes in surface temperature. *Journal of Climate* 30(6):2251–2267.
- Conn, P.B., Chernook, V.I., Moreland, E.E., et al. 2021. Aerial survey estimates of polar bears and their tracks in the Chukchi Sea. *PLoS ONE* 16(5): e0251130. <https://doi.org/10.1371/journal.pone.0251130>.
- Cooke, J.G. 2018a. *Balaenoptera musculus* (errata version published in 2019). *The IUCN Red List of Threatened Species* 2018: e.T2477A156923585.
- Cooke, J.G. 2018b. *Balaenoptera physalus*. *The IUCN Red List of Threatened Species* 2018: e.T2478A50349982.
- Cooke, J.G. and Zerbini, A.N. 2018. *Eubalaena australis*. *The IUCN Red List of Threatened Species* 2018: e.T8153A50354147. <https://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T8153A50354147.en>.

- Cooke, J.G., Zerbini, A.N. & Taylor, B.L. 2018. *Balaenoptera bonaerensis*. *The IUCN Red List of Threatened Species* 2018: e.T2480A50350661. <https://dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T2480A50350661.en>.
- Cornish, S.B., Johnson, H.L., Mallet, R.D.C., et al. 2022. Rise and fall of sea ice production in the Arctic Ocean's ice factories. *Nature Communications* 13:7800.
- COSEWIC. 2018. COSEWIC assessment and status report on the Polar Bear *Ursus maritimus* in Canada. *Committee on the Status of Endangered Wildlife in Canada*. Ottawa.
- COSEWIC. 2020. COSEWIC assessment and status report on the Beluga Whale *Delphinapterus leucas*, Eastern High Arctic - Baffin Bay population, Cumberland Sound population, Ungava Bay population, Western Hudson Bay population, Eastern Hudson Bay population and James Bay population in Canada. *Committee on the Status of Endangered Wildlife in Canada*. Ottawa. <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/beluga-whale-2020.html>.
- Crawford, J.A., Quakenbush, L.T. and Citta, J.J. 2015. A comparison of ringed and bearded seal diet, condition and productivity between historical (1975–1984) and recent (2003–2012) periods in the Alaskan Bering and Chukchi seas. *Progress in Oceanography* 136:133–150.
- Crockford, S. J. 2002. Animal domestication and heterochronic speciation: the role of thyroid hormone. pg. 122–153. In: N. Minugh-Purvis and K. McNamara (eds.) *Human Evolution Through Developmental Change*. Baltimore, Johns Hopkins University Press.
- Crockford, S. J. 2003. Thyroid hormone phenotypes and hominid evolution: a new paradigm implicates pulsatile thyroid hormone secretion in speciation and adaptation changes. *International Journal of Comparative Biochemistry and Physiology Part A* 135(1):105–129.
- Crockford, S. J. 2004. Animal Domestication and Vertebrate Speciation: A Paradigm for the Origin of Species. Ph.D. dissertation. University of Victoria, Canada.
- Crockford, S. J. 2006. *Rhythms of Life: Thyroid Hormone and the Origin of Species*. Trafford, Victoria.
- Crockford, S. J. 2008. Be careful what you ask for: archaeozoological evidence of mid-Holocene climate change in the Bering Sea and implications for the origins of Arctic Thule. In: *Islands of Inquiry: Colonisation, Seafaring and the Archaeology of Maritime Landscapes*, 113–131. G. Clark, F. Leach and S. O'Connor (eds.). Terra Australis 29 ANU EPress, Canberra.
- Crockford, S. J. 2009. Evolutionary roots of iodine and thyroid hormones in cell-cell signaling. *Integrative and Comparative Biology* 49:155–166.
- Crockford, S. J. 2012. A History of Polar Bears, Ringed Seals, and other Arctic and North Pacific Marine Mammals over the Last 200,000 Years. Report prepared for The State of Alaska Department of Commerce, Community and Economic Development and The University of Alaska Fairbanks. Pacific Identifications Inc., Victoria, British Columbia.
- Crockford, S. J. 2014a. On the beach: walrus haulouts are nothing new. Global Warming Policy Foundation Briefing Paper 11, London. <http://www.thegwpf.org/susan-crockford-on-the-beach-2/>.
- Crockford, S. J. 2014b. The walrus fuss: walrus haulouts are nothing new. Global Warming Policy Forum. http://www.thegwpf.org/gwpftv/?tubepress_item=cwaAwsS2O0Y&tubepress_page=2.
- Crockford, S.J. 2015a. The Arctic fallacy: sea ice stability and the polar bear. Global Warming Policy Foundation Briefing Paper 16, London.
- Crockford, S.J. 2015b. A harrowing encounter: polar bear vs. lawyer in Labrador. *Range Magazine* Spring: 42–43.
- Crockford, S.J. 2017. Testing the hypothesis that routine sea ice coverage of 3–5 mkm² results in a greater than 30% decline in population size of polar bears (*Ursus maritimus*). *PeerJ Preprints*, 2 March 2017. Doi: 10.7287/peerj.preprints.2737v3.

- Crockford, S.J. 2018a. Prehistoric mountain goat *Oreamnos americanus* mother lode near Prince Rupert, British Columbia and implications for the manufacture of high-status ceremonial goods. *Journal of Island and Coastal Archaeology* 13(3):319–340.
- Crockford, S.J. 2018b. State of the Polar Bear Report 2017. Global Warming Policy Foundation Report 29, London.
- Crockford, S.J. 2018c. 'Polar bears keep thriving even as global warming alarmists keep pretending they're dying.' *National Post/Financial Post* (Canada), 27 February. <http://business.financialpost.com/opinion/polar-bears-keep-thriving-even-as-global-warming-alarmists-keep-pretending-theyre-dying>.
- Crockford, S.J. 2018d. 'The real story behind the famous starving polar bear video reveals more manipulation.' *National Post/Financial Post* (Canada), 29 August. <https://business.financialpost.com/opinion/the-real-story-behind-the-famous-starving-polar-bear-video-reveals-more-manipulation>.
- Crockford, S.J. 2018e. White lie: the cruel abuse of a starving polar bear. August. Global Warming Policy Forum, London. <https://www.youtube.com/watch?v=Z7KTfPICrgY>.
- Crockford, S.J. 2019a. *State of the Polar Bear 2018*. Global Warming Policy Foundation Report 32, London.
- Crockford, S.J. 2019b. 'Netflix is lying about those falling walruses. It's another 'tragedy porn' climate hoax.' *National Post/Financial Post* (Canada), 24 April. <https://business.financialpost.com/opinion/netflix-is-lying-about-those-falling-walruses-its-another-tragedy-porn-climate-hoax>.
- Crockford, S.J. 2019c. 'Netflix, Attenborough and cliff-falling walruses: the making of a false climate icon.' May. Global Warming Policy Forum, London. <https://www.youtube.com/watch?v=latVKZZcPG0>.
- Crockford, S.J. 2019d. 'The truth about Attenborough's falling walruses'. September. Global Warming Policy Foundation, London. <https://www.youtube.com/watch?v=tFcwAKZEnHY>.
- Crockford, S.J. 2019e. 'No climate emergency for polar bears.' September. Global Warming Policy Foundation, London. <https://www.youtube.com/watch?v=jQRle6pgBCY>.
- Crockford, S.J. 2019f. 'Falling walrus: Attenborough tacitly admits Netflix deception.' December. Global Warming Policy Foundation, London. <https://www.youtube.com/watch?v=U5Ji6ME3Vlo>.
- Crockford, S.J. 2020a. 'New footage reveals Netflix faked walrus climate deaths.' 19 November. Global Warming Policy Forum, London. <https://www.youtube.com/watch?v=kV8d26oziVM>.
- Crockford, S.J. 2020b. *State of the Polar Bear 2019*. Global Warming Policy Foundation Report 39, London.
- Crockford, S.J. 2021. *The State of the Polar Bear Report 2020*. Global Warming Policy Foundation Report 48, London.
- Crockford, S.J. 2022a. *The State of the Polar Bear Report 2021*. Global Warming Policy Foundation Note 29, London.
- Crockford, S.J. 2022b. Polar bear fossil and archaeological records from the Pleistocene and Holocene in relation to sea ice extent and open water polynyas. *Open Quaternary* 8(1):7.
- Crockford, S.J. 2023. 'Opinion: What climate alarmism about polar bears gets wrong.' *National Post/Financial Post* (Canada), 13 January. <https://financialpost.com/opinion/what-climate-alarmism-about-polar-bears-gets-wrong>.
- Crockford, S., and Frederick, G. 2007. Sea ice expansion in the Bering Sea during the Neoglacial: evidence from archaeozoology. *The Holocene* 17:699–706.
- Crockford, S.J. and Frederick, G. 2011. Neoglacial sea ice and life history flexibility in ringed and fur seals. pg. 65–91. In: T. Braje and R. Torrey (eds). *Human and Marine Ecosystems: Archaeology and Historical Ecology of Northeastern Pacific Seals, Sea Lions, and Sea Otters*. U. California Press.
- Crockford, S.J. and Geist, V. 2018. Conservation Fiasco. *Range Magazine*, Winter 2017/2018, pg. 26–27.

- De Augustinis, F. 2022a. 'Fish-feed industry turns to krill, with unknown effects on the Antarctic ecosystem.' *Mongabay News*, 19 October. <https://news.mongabay.com/2022/10/fish-feed-industry-turns-to-krill-with-unknown-effects-on-the-antarctic-ecosystem/>.
- De Augustinis, F. 2022b. 'Negotiations to conserve Antarctic Ocean in stalemate on many issues.' *Mongabay News*, 7 November. <https://news.mongabay.com/2022/11/negotiations-to-conserve-antarctic-ocean-end-in-stalemate-on-many-issues/>.
- de Bruyn, P.J.N., Tosh, C.A. and Terauds, A. 2013. Killer whale ecotypes: is there a global model? *Biological Reviews* 88(1):62–80.
- DFO Canada. 2022. Report on the Progress of Recovery Strategy Implementation for the Beluga Whale (*Delphinapterus leucas*) (DFO), St. Lawrence Estuary Population in Canada, for the Period 2012 to 2019. *Species at Risk Act Recovery Strategy Report Series*. Fisheries and Oceans Canada, Ottawa. <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/report-progress-recovery-document/beluga-whale-2022.html>.
- Dickie, G. 2022a. 'In Antarctica, does a burgeoning krill fishery threaten wildlife?' *Reuters*, 24 February. <https://www.reuters.com/business/cop/antarctica-does-burgeoning-krill-fishery-threaten-wildlife-2022-02-24/>.
- Dickie, G. 2022b. 'Canada's Hudson Bay polar bear population plummets as climate change warms Arctic.' *Reuters*, 23 December. <https://www.reuters.com/world/americas/canadas-hudson-bay-polar-bear-population-plummets-climate-change-warms-arctic-2022-12-23/>.
- Doherty, S. 2016. 'Common murre update: growing awareness of sea bird die-off thanks to citizen reporting.' *Alaska Fish & Wildlife News*, April. https://www.adfg.alaska.gov/index.cfm?adfg=wildlifeneews.view_article&articles_id=770.
- Doran, T. 2020. Emails from the edge: Svalbard's polar bears are sending messages to scientists. *CNN*, 9 Dec 2020. <https://www.cnn.com/2020/12/09/europe/polar-bears-svalbard-jon-aars-norwegian-polar-institute-spc-intl-c2e/index.html>.
- Dunmall, K.M., McNicholl, D.G., Zimmerman, C.E., et al. 2022. First juvenile chum salmon confirms successful reproduction for Pacific salmon in the North American Arctic. *Canadian Journal of Fisheries and Aquatic Sciences* 79(5):703–707.
- Durban, J.W., Fearnbach, H., Burrows, D.G. et al. 2017. Morphological and ecological evidence for two sympatric forms of Type B killer whale around the Antarctic Peninsula. *Polar Biology* 40:231–236.
- Durban, J.W., Fearnbach, H., Paredes, A., et al. 2021. Size and body condition of sympatric killer whale ecotypes around the Antarctic Peninsula. *Marine Ecology Progress Series* 677:207–217.
- Durner, G.M., Laidre, K.L., and York, G.S. (eds). 2018. Polar Bears: Proceedings of the 18th Working Meeting of the IUCN/SSC Polar Bear Specialist Group, 7–11 June 2016, Anchorage, Alaska. Gland, Switzerland and Cambridge, UK: IUCN.
- Dyck, M., Lukacs, P.M. and Ware, J.V. 2021a. Recovery from reduction: M'Clintock Channel polar bear subpopulation, Nunavut, Canada. *Arctic* 74(4):509–524.
- Dyck, M., Dunham, K.D., Ware, J.V., et al. 2021b. Re-estimating the abundance of the Davis Strait polar bear subpopulation by genetic mark-recapture. Final Report [amended 9 May 2022], Government of Nunavut, Department of Environment, Iglulik.
- Dyck, M., Regehr, E.V. and Ware, J.V. 2023. Demographic assessment using physical and genetic sampling finds stable polar bear subpopulation in Gulf of Boothia, Canada. *Marine Mammal Science* 39:151–174.
- Environment and Climate Change Canada 2021. *Species at Risk in Nunavut 2021*. Iqaluit, NU. <https://www.canada.ca/en/environment-climate-change/services/species-risk-education-centre/species-risk-nunavut-2021>.

Farmer, J.R., Sigman, D.M., Granger, J., et al. 2021. Arctic Ocean stratification set by sea level and freshwater inputs since the last ice age. *Nature Geoscience* 14:684–689. <https://doi.org/10.1038/s41561-021-00789-y>.

Fearnbach, H., Durban, J.W., Ellifrit, D.K. et al. 2019. Abundance of type A killer whales (*Orcinus orca*) in the coastal waters off the western Antarctic Peninsula. *Polar Biology* 42:1477–1488.

Fearnbach, H., Durban, J.W., Ellifrit, D.K. et al. 2021. A decade of photo-identification reveals contrasting abundance and trends of Type B killer whales in the coastal waters of the Antarctic Peninsula. *Marine Mammal Science* 38(1):58–72.

Fedewa, E.J., Jackson, T.M., Richar, J.I., et al. 2020. Recent shifts in northern Bering Sea snow crab (*Chionoecetes opilio*) size structure and the potential role of climate-mediated range contraction. *Deep Sea Research Part II: Topical Studies in Oceanography* 181–182:1–4878.

Fischbach, A.S., Taylor, R.L. and Jay, C.V. 2022. Regional walrus abundance estimate in the United States Chukchi Sea in autumn. *Journal of Wildlife Management* 86:e22256.

Fogt, R.L., Sleinkofer, A.M., Raphael, M.N. and Handcock, M.S. 2022. A regime shift in seasonal total Antarctic sea ice extent in the twentieth century. *Nature Climate Change* 12:54–62.

Foote, A.D., Martin, M.D., Louis, M., et al. 2019. Killer whale genomes reveal a complex history of recurrent admixture and vicariance. *Molecular Ecology* 28(14):3427–3444.

Fretwell, P.T. and Trathan, P.N. 2019. Emperors on thin ice: three years of breeding failure at Halley Bay. *Antarctic Science* 31(3):133–138.

Fretwell, P.T. and Trathan, P.N. 2020. Discovery of new colonies by Sentinel2 reveals good and bad news for emperor penguins. *Remote Sensing in Ecology and Conservation* 7(2): 139–153.

Fretwell, P.T., LaRue, M.A., Morin, P., Kooyman, G.L., Wienecke, B., Ratcliffe, N., Fox, A.J., Fleming, A.H., Porter, C. and Trathan, P.N. 2012. An emperor penguin population estimate: the first global, synoptic survey of a species from space. *PLoS One* 7: e33751.

Frey, K.E., Comiso, J.C., Cooper, L.W., et al. 2022. Arctic Ocean primary productivity: the response of marine algae to climate warming and sea ice decline. In: *2022 Arctic Report Card*, NOAA. <https://doi.org/10.25923/0je1-te61>.

Gamillo, E. 2022. 'Watch thousands of beluga whales migrate from the Arctic.' *Smithsonian Magazine*, 15 July. <https://www.smithsonianmag.com/smart-news/watch-thousands-of-beluga-whales-migrate-to-a-bay-in-canada-180980425/>.

George, J.C., Moore, S.E. and Thewissen, J.G.M. 2020. Bowhead whales: recent insights into their biology, status, and resilience. *2020 Arctic Report Card*, NOAA. DOI: 10.25923/cppm-n265.

George, J. 2022. 'Nunavut hunters cry foul over new polar bear management scheme.' *CBC News*, 16 June. <https://www.cbc.ca/news/canada/north/polar-bear-management-gulf-of-boothia-1.6489961>.

Gill, V. 2022. 'Canada's polar-bear capital Churchill warms too fast for bears.' *BBC News*, 18 December. <https://www.bbc.com/news/science-environment-63603489>.

Government of Canada 2022. 'Inuit polar bear experts share their knowledge in a ground-breaking new study'. *Government of Canada Press release*, 12 August. <https://www.canada.ca/en/polar-knowledge/polar-blog-articles/inuit-polar-bear-experts-share-their-knowledge-in-a-ground-breaking-new-study.html>.

Haavik, E. 2022. 'Svalbard's polar bears persist as sea ice melts — but not forever. *The World*, 21 July. <https://theworld.org/stories/2022-07-21/svalbard-s-polar-bears-persist-sea-ice-melts-not-forever>.

Halliday, W., McBeth, S., Vergara, V., et al. 2022, Marine mammals in changing Arctic Ocean. *Polar Knowledge: Aqhaliat Report*, Volume 4, Polar Knowledge Canada, p. 58–83. DOI: 10.35298/pkc.2021.03.eng.

Hamamdjian, D. and Jones, A.M. 2022. 'New polar bear cubs a good sign, but species still in danger, experts say.' *CTV News*, 8 May. <https://www.ctvnews.ca/climate-and-environment/new-polar-bear-cubs-a-good-sign-but-species-still-in-danger-experts-say-1.5894379>.

Hamilton, S.G. and Derocher, A.E. 2019. Assessment of global polar bear abundance and vulnerability. *Animal Conservation* 22:83–95.

Hausfather, Z. and Peters, G.P. 2020. Emissions – the 'business as usual' story is misleading ["Stop using the worst-case scenario for climate warming as the most likely outcome — more-realistic baselines make for better policy"]. *Nature* 577: 618–620.

Heide-Jørgensen, M.P., Chambault, P., Jansen, T. et al. 2022. A regime shift in the Southeast Greenland marine ecosystem. *Global Change Biology* 29(3):668–685.

Herr, H., Viquerat, S., Devas, F., et al. 2022. Return of large fin whale feeding aggregations to historical whaling grounds in the Southern Ocean. *Scientific Reports* 12:9458.

Herreman, J. and Peacock, E. 2013. Polar bear use of a persistent food subsidy: Insights from non-invasive genetic sampling in Alaska. *Ursus* 24(2):148–163.

Holt, R.D. 2019. Pacific walrus terrestrial haul-out use on Togiak National Wildlife Refuge, Alaska, 1985–2018. US Fish and Wildlife Service Report, Togiak National Wildlife Refuge. Dillingham, Alaska.

Hückstädt, L. 2015a. *Leptonychotes weddellii*. *The IUCN Red List of Threatened Species* 2015: e.T11696A45226713. <https://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T11696A45226713.en>.

Hückstädt, L. 2015b. *Lobodon carcinophaga*. *The IUCN Red List of Threatened Species* 2015: e.T12246A45226918.

Hückstädt, L. 2015c. *Hydrurga leptonyx*. *The IUCN Red List of Threatened Species* 2015: e.T10340A45226422.

Hückstädt, L. 2015d. *Ommatophoca rossii*. *The IUCN Red List of Threatened Species* 2015: e.T15269A45228952.

Innes, S., Heide-Jørgensen, M.P., Laake, J.L., Laidre, K.L., Cleator, H.J., Richard, P., and Stewart, R.E.A. 2002. Surveys of belugas and narwhals in the Canadian High Arctic in 1996. *NAMMCO Scientific Publications* 4:169–190.

IWC. 2016. Report of the Scientific Committee of the International Whaling Commission 2016: Annex H: Report of the Sub-Committee on Other Southern Hemisphere Whale Stocks 44 (International Whaling Commission, Bled, Slovenia).

IWC 2018a; Blue whale. *International Whaling Commission*, <https://iwc.int/about-whales/whale-species/blue-whale>.

IWC. 2018b. Fin whale. *International Whaling Commission*, <https://iwc.int/about-whales/whale-species/fin-whale>.

Jacobo, J. 2022. 'More interactions between humans and polar bears are likely as sea ice melts due to climate change, scientists say'. *ABC News*, 10 November. <https://abcnews.go.com/International/interactions-humans-polar-bears-sea-ice-melts-due/story?id=93004806>.

Jenouvrier, S., Caswell, H., Barbraud, C., Holland, M., Stroeve, J. and Weimerskirch, H. 2009. Demographic models and IPCC climate projections predict the decline of an emperor penguin population. *Proceedings of the National Academy of Science USA* 106: 1844–1847.

Jenouvrier, S., Holland, M., Iles, D., et al. 2020. The Paris Agreement objectives will likely halt future declines of emperor penguins. *Global Change Biology* 26(3): 1170–1184.

Kaler, R., Sheffield, G., Backensto, S., et al. 2022. Partnering in search of answers: seabird die-offs in the Bering and Chukchi Seas. *NOAA Arctic Report Card 2022*. <https://arctic.noaa.gov/Report-Card/Report-Card-2022/ArtMID/8054/ArticleID/1006/Partnering-in-Search-of-Answers-Seabird-Die-offs-in-the-Bering-and-Chukchi-Seas>.

- Kaufman, M. 2019. Village of the Sea Bears: Churchill's famed polar bear population is plunging. *Mashable*, 20 November. <https://mashable.com/feature/polar-bears-churchill-population-decline/>.
- Kovacs, K.M. 2016. *Odobenus rosmarus ssp. rosmarus*. The IUCN Red List of Threatened Species 2016: e.T15108A66992323. <http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T15108A66992323.en>.
- Krüger, L., Huerta, M. F., Santa Cruz, F., and Cárdenas, C. A. 2020. Antarctic krill fishery effects over penguin populations under adverse climate conditions: Implications for the management of fishing practices. *Ambio* 50(3):560–571.
- Labrousse, S., Williams, G., Tamura, T. et al. 2018. Coastal polynyas: Winter oases for subadult southern elephant seals in East Antarctica. *Scientific Reports* 8:3183.
- Lang, A., Yang, S. and Kaas, E. 2017. Sea ice thickness and recent Arctic warming. *Geophysical Research Letters* 44:409–418. <https://doi.org/10.1002/2016GL071274>.
- LaRue, M., Salas, L., Nur, N., et al. 2021. Insights from the first global population estimate of Weddell seals in Antarctica. *Science Advances* 7:eabh3674.
- Leathlobhair, M.N., Perri, A.R., Irving-Pease, E.K., et al. [including Crockford, S.J.]. 2018. The evolutionary history of dogs in the Americas. *Science* 361:81–85.
- Lewis, K. M., van Dijken, G.L. and Arrigo, K.R. 2020. Changes in phytoplankton concentration now drive increased Arctic Ocean primary production. *Science* 369(6500):198–202.
- Lillie, K.M., Gese, E.M., Atwood, T.C., et al. 2019. Use of subsistence-harvested whale carcasses by polar bears in the Southern Beaufort Sea. *Arctic* 72(4):404–412.
- Lippold, A., Bourgeon, S., Aars, J., et al. 2019. Temporal trends of persistent organic pollutants in Barents Sea polar bears (*Ursus maritimus*) in relation to changes in feeding habits and body condition. *Environmental Science and Technology* 53(2):984–995.
- Lowther, A.D., Staniland, I., Lydersen, C., and Kovacs, K.M. 2020. Male Antarctic fur seals: neglected food competitors of bioindicator species in the context of an increasing Antarctic krill fishery. *Scientific Reports* 10(1):18436.
- MacCracken, J.G., Beatty, W.S., Garlich-Miller, J.L., et al. 2017. *Final Species Status Assessment for the Pacific walrus (Odobenus rosmarus divergens)*. US Fish and Wildlife Service, Anchorage, Alaska.
- McBride, M.M., Schram Stokke, O., Renner, A.H.H., et al. 2021. Antarctic krill *Euphausia superba*: spatial distribution, abundance, and management of fisheries in a changing climate. *Marine Ecology Progress Series* 668:185–214.
- McKechnie, I., Moss, M.L. and Crockford, S.J. 2020. Domestic dogs and wild canids on the Northwest Coast of North America: animal husbandry in a region without agriculture? *Journal of Anthropological Archaeology* 60:101209.
- Mangerud, J. and Svendsen, J.I. 2018. The Holocene Thermal Maximum around Svalbard, Arctic North Atlantic; molluscs show early and exceptional warmth. *The Holocene* 28: 65–83.
- March, D., Drago, M., Gazo, M. et al. 2021. Winter distribution of juvenile and sub-adult male Antarctic fur seals (*Arctocephalus gazella*) along the western Antarctic Peninsula. *Scientific Reports* 11:22234.
- Marine Mammal Science. 2023a. *Leptonychotes weddellii* (Weddell seal). Marine Mammal Science Education Committee. <https://marinemammalscience.org/facts/leptonychotes-weddellii/>.
- Marine Mammal Science. 2023b. *Lobodon carcinophaga* (Crabeater seal). Marine Mammal Science Education, <https://marinemammalscience.org/facts/lobodon-carcinophaga/>.
- Martinez-Levasseur, L.M., Furgal, C.M., Hammill, M.O., et al. 2021. New migration and distribution patterns of Atlantic walruses (*Odobenus rosmarus rosmarus*) around Nunavik (Québec, Canada) identified using Inuit Knowledge. *Polar Biology* 44:1833–1845.
- Matishov, G.G., Chelintsev, N.G., Goryaev, Yu. I., et al. 2014. Assessment of the amount of polar bears (*Ursus maritimus*) on the basis of perennial vessel counts. *Doklady Earth Sciences* 458 (2):1312–1316.

Meier, W. 2019. September monthly mean extent and trends for 1979–2019, showing overall trend and trends for the most recent 13 years, and the steepest 13 years in the 41-year record. Figure 3b, in 'Falling Up' [sea ice conditions for September 2019], *NSIDC Arctic Sea Ice News & Analysis*, 3 October. <http://nsidc.org/arcticseaicenews/2019/10/falling-up/>.

Meier, W.N. and Stewart, J.S. 2019. Assessing uncertainties in sea ice extent climate indicators. *Environmental Research Letters* 14:035005. <https://doi.org/10.1088/1748-9326/aaf52c>.

Meier, W., Petty, A., Hendricks, S., et al. 2022. Sea Ice. In: *2021 Arctic Report Card*, NOAA. <https://doi.org/10.25923/xyp2-vz45>.

Meyer, B., Atkinson, A., Bernard, K.S. et al. 2020. Successful ecosystem-based management of Antarctic krill should address uncertainties in krill recruitment, behaviour and ecological adaptation. *Nature Communications Earth & Environment* 1:28.

Miller, E.N., Lunn, N.J., McGeachy, D., and Derocher, A.E. 2022. Autumn migration phenology of polar bears (*Ursus maritimus*) in Hudson Bay, Canada. *Polar Biology* 45:1023–1034.

Miller, S., Schliebe, S. and Proffitt, K. 2006. Demographics and behavior of polar bears feeding on bowhead whale carcasses at Barter and Cross Islands, Alaska, 2002–2004. Alaska Outer Continental Shelf (OCS) Study MMS 2006–14, US Dept. of the Interior, Minerals Management Service, Anchorage.

Miller, S., Wilder, J. and Wilson, R.R. 2015. Polar bear-grizzly bear interactions during the autumn open-water period in Alaska. *Journal of Mammalogy* 96(6):1317–1325.

Molnár, P.K., Bitz, C.M., Holland, M.M., et al. 2020. Fasting season length sets temporal limits for global polar bear persistence. *Nature Climate Change* 10:732–738.

Molnár, P.K., Derocher, A.E., Theimann, G., and Lewis, M.A. 2010. Predicting survival, reproduction and abundance of polar bears under climate change. *Biological Conservation* 143:1612–1622.

Naishadham, S. 2022. 'Canadian polar bears near 'bear capital of the world' dying at fast rate.' Associated Press, 22 December. <https://apnews.com/article/science-arctic-polar-bears-hudson-bay-climate-and-environment-efd19bb16967dd058b2ea39f9d676dd0>.

NAMMCO. 2021. Atlantic walrus. *North Atlantic Marine Mammal Commission Report for 2016*, updated January 2021. Tromsø, Norway. <https://nammco.no/topics/atlantic-walrus/>.

Norwegian Polar Institute (NPI). 2022a. Condition in adult polar bear males. Environmental monitoring of Svalbard and Jan Mayen (MOSJ). <http://www.mosj.no/en/fauna/marine/polar-bear.html>.

Norwegian Polar Institute (NPI). 2022b. Polar bear cubs per litter. Environmental monitoring of Svalbard and Jan Mayen (MOSJ). <http://www.mosj.no/en/fauna/marine/polar-bear.html>.

Norwegian Polar Institute (NPI). 2022c. Production of polar bear cubs. Environmental monitoring of Svalbard and Jan Mayen (MOSJ). <http://www.mosj.no/en/fauna/marine/polar-bear.html>.

Norwegian Polar Institute (NPI). 2022d. Walrus population in Svalbard. *Environmental Monitoring of Svalbard and Jan Mayen* (MOSJ). <http://www.mosj.no/en/fauna/marine/walrus-population.html>.

NSIDC 2014. '2014 melt season in review'. *US National Snow and Ice Data Center*, 7 October. <https://nsidc.org/arcticseaicenews/2014/10/2014-melt-season-in-review/>.

NSIDC 2021. 'September Turning: Antarctica's unusual sea ice maximum'. *US National Snow and Ice Data Center*, 5 October. <http://nsidc.org/arcticseaicenews/2021/10/september-turning/>.

NSIDC 2022. 'Ain't no sunshine when she's gone'. *US National Snow and Ice Data Center*, 4 October. <https://nsidc.org/arcticseaicenews/2022/10/no-sunshine-when-shes-gone/>.

NSIDC 2023. 'December lows'. *US National Snow and Ice Data Center*, 5 January. <http://nsidc.org/arcticseaicenews/2023/01/december-lows/>.

Obbard, M.E., Theimann, G.W., Peacock, E., et al. (eds.) 2010. Polar Bears: Proceedings of the 15th meeting of the Polar Bear Specialist Group IUCN/SSC, 29 June–3 July, 2009, Copenhagen, Denmark. Gland, Switzerland and Cambridge UK, IUCN.

- Ogilvie, A.E.J., Barlow, L.K. and Jennings, A.E. 2000. North Atlantic climate c. AD 1000: Millennial reflections on the Viking discoveries of Iceland, Greenland and North America. *Weather* 55: 34–45.
- Omrani, N.-E., Keenlyside, N., Matthes, K., et al. 2022. Coupled stratosphere-troposphere-Atlantic multidecadal oscillation and its importance for near-future climate projection. *Climate and Atmospheric Science* 5:59.
- PBSG. 2019. 'Status Report on the World's Polar Bear Subpopulations July 2019'. *IUCN/SSC Polar Bear Specialist Group*, 20 September.
- PBSG. 2021. 'Status Report on the World's Polar Bear Subpopulations at 31 July 2021'. *IUCN Polar Bear Specialist Group*, 3 December. <https://www.iucn-pbsg.org/>.
- Pearce, J.M., Dooley, J., Patil, V., et al., 2022. Arctic geese of North America. *NOAA Arctic Report Card 2022*. <https://arctic.noaa.gov/Report-Card/Report-Card-2022/ArtMID/8054/ArticleID/1000/Arctic-Geese-of-North-America>.
- Perrin, W.F. and Brownell Jr, R.L. 2009. Minke whales *Balaenoptera acutorostrata* and *B. bonaerensis* in *Encyclopedia of Marine Mammals* (eds W. Perrin, B. Wursig, & J.G.M. Thewissen), pg. 733–735. Elsevier.
- Pinkerton, M.H., Boyd, P.W., Deppeler, S., et al. 2021. Evidence for the impact of climate change on primary producers in the Southern Ocean. *Frontiers in Ecology and Evolution* 9:592027.
- Piñones, A. and Fedorov, A.V. 2016. Projected changes of Antarctic krill habitat by the end of the 21st century. *Geophysical Research Letters* 43(16): 8580–8589.
- Pitman, R.L. 2011. Antarctic killer whale: top of the food chain at the bottom of the world. *Whale-watcher aka Journal of the American Cetacean Society* 40(1):39–45.
- Pitman, R.L., Perryman, W.L., DeRoi, D. and Eilers, E. 2007. A dwarf form of killer whale in Antarctica. *Journal of Mammalogy* 88(1):43–48.
- Purser, A., Hehemann, L., Boehringer, L., et al. 2022. A vast icefish breeding colony discovered in the Antarctic. *Current Biology*, 32(4): 842–850.
- Rackow, T., Danilov, S., Goessling, H.F. et al. 2022. Delayed Antarctic sea-ice decline in high-resolution climate change simulations. *Nature Communications* 13:637.
- Ramirez, R. 2022. 'Billions of snow crabs have disappeared from the waters around Alaska. Scientists say overfishing is not the cause.' *CNN*, 16 October. <https://www.cnn.com/2022/10/16/us/alaska-snow-crab-harvest-canceled-climate/index.html>.
- Regehr, E.V., Laidre, K.L., Akçakaya, H.R., et al. 2016. Conservation status of polar bears (*Ursus maritimus*) in relation to projected sea-ice declines. *Biology Letters* 12: 20160556.
- Regehr, E.V., Hostetter, N.J., Wilson, R.R., et al. 2018. Integrated population modeling provides the first empirical estimates of vital rates and abundance for polar bears in the Chukchi Sea. *Scientific Reports* 8(1). <https://doi.org/10.1038/s41598-018-34824-7>.
- Richardson, E. 2019. Case study: Polar bears and the decline of sea ice. In: State of Canada's Arctic Seas, Niemi, A., Ferguson, S., Hedges, K., et al. (eds.). Canadian Technical Report Fisheries and Aquatic Sciences 3344.
- Risch, D., Norris, T., Curnock, M. and Freidlaender, A. 2019. Common and Antarctic minke whales: conservation status and future research directions. *Frontiers in Marine Science* 6:247.
- Rode, K.D., Wilson, R.R., Douglas, D.C., et al. 2018. Spring fasting behavior in a marine apex predator provides an index of ecosystem productivity. *Global Change Biology* 24(1):410–423.
- Rode, K. D., Regehr, E.V., Bromaghin, J. F., et al. 2021. Seal body condition and atmospheric circulation patterns influence polar bear body condition, recruitment, and feeding ecology in the Chukchi Sea. *Global Change Biology* 27:2684–2701. <https://doi.org/10.1111/gcb.15572>.
- Rode, K.D., Regehr, E.V., Douglas, D., et al. 2014. Variation in the response of an Arctic top predator experiencing habitat loss: feeding and reproductive ecology of two polar bear populations. *Global Change Biology* 20(1):76–88.

- Rode, K.D., Wilson, R.R., Regehr, E.V., et al. 2015. Increased land use by Chukchi Sea polar bears in relation to changing sea ice conditions. *PLoS ONE* 10(11):e0142213.
- Rogers, M.C., Peacock, E., Simac, K., et al. 2015. Diet of female polar bears in the southern Beaufort Sea of Alaska: evidence for an emerging alternative foraging strategy in response to environmental change. *Polar Biology* 38(7):1035–1047.
- Ryan, J. 2022. 'Antarctica's fin whale population is rebounding. Here's why that's a big deal.' *CNET Science*, 12 July. <https://www.cnet.com/science/biology/antarcticas-fin-whale-population-is-rebounding-heres-why-thats-a-big-deal/>.
- Scott, M. 2020. 'Understanding climate: Antarctic sea ice extent.' *NOAA Climate News Feature*, 28 April. <https://www.climate.gov/news-features/understanding-climate/understanding-climate-antarctic-sea-ice-extent>.
- Shiozaki, T., Fukiwara, A., Sugie, K., et al. 2022. Bottom-associated phytoplankton bloom and its expansion in the Arctic Ocean. *Global Change Biology* 28:7286–7295.
- Siddon, E. 2021. Ecosystem Status Report for the Eastern Bering Sea, *Stock Assessment and Fishery Evaluation Report*. North Pacific Fishery Management Council, Anchorage, AK.
- Siegel, V. and Watkins, J.L. 2016. Distribution, biomass and demography of Antarctic krill, *Euphausia superba*. In: Siegel, V. (ed.), *Biology and Ecology of Antarctic Krill*, pp 21–100. Advances in Polar Ecology. Springer.
- Smith, Jr., W.O. 2022. Primary productivity measurements in the Ross Sea, Antarctica: a regional synthesis. *Earth Systems Science Data* 14:2737–2747.
- Smith, T.G. and Martin, A.R. 1994. Distribution and movements of belugas, *Delphinapterus leucas*, in the Canadian High Arctic. *Canadian Journal of Fisheries and Aquatic Sciences* 51:1653–1663.
- Smith, T.S. 2022. Human garbage is a plentiful but dangerous source of food for polar bears finding it harder to hunt seals on dwindling sea ice. *The Conversation*, 20 July. <https://theconversation.com/human-garbage-is-a-plentiful-but-dangerous-source-of-food-for-polar-bears-finding-it-harder-to-hunt-seals-on-dwindling-sea-ice-183968>.
- Smith, T.S., Derocher, A.E., Mazur, R.L., et al. 2022. Anthropogenic food: an emerging threat to polar bears. *Oryx*, in press. <https://doi.org/10.1017/S0030605322000278>.
- Snoeijs-Leijonmalm, P., Flores, H., Sakinan, S., et al. 2022. 'Unexpected fish and squid in the central Arctic deep scattering layer. *Science Advances* 8(7):eabj7536.
- Snook, J. 2022. 'How blending Inuit knowledge and western science has helped improve polar bear health — and why a trade ban would hurt.' *The Conversation*, 18 April. <https://theconversation.com/how-blending-inuit-knowledge-and-western-science-has-helped-improve-polar-bear-health-and-why-a-trade-ban-would-hurt-173579>.
- Southwell, C., Emmerson, L., Takahashi, A., et al. 2017. Large-scale population assessment informs conservation management for seabirds in Antarctica and the Southern Ocean: A case study of Adelie penguins. *Global Ecology and Conservation* 9: 104–115.
- Stirling, I. and Derocher, A.E. 2012. Effects of climate warming on polar bears: a review of the evidence. *Global Change Biology* 18:2694–2706 doi:10.1111/j.1365–2486.2012.02753.x.
- Stirling and Kiliaan. 1980. Population ecology studies of the polar bear in northern Labrador. Canadian Wildlife Service Occasional Paper No. 42.
- Stroeve, J., Holland, M.M., Meier, W., et al. 2007. Arctic sea ice decline: Faster than forecast. *Geophysical Research Letters* 34:L09501.
- Stroeve, J., Markus, T., Boisvert, L., et al. 2014. Changes in Arctic melt season and implications for sea ice loss. *Geophysical Research Letters* 41:1216–1224.
- Strycker, N. 2020. 'Antarctica's most numerous penguin has suffered huge declines, expedition finds.' *Audubon Magazine*, 10 February. <https://www.audubon.org/news/antarcticas-most-numerous-penguin-has-suffered-huge-declines-expedition-finds>.

- Strycker, N. and Lynch, H. 2022. *Pygoscelis antarcticus* (Green Status assessment). *The IUCN Red List of Threatened Species* 2022: e.T22697761A2269776120221. Accessed on 30 January 2023.
- Strycker, N., Wethington, M., Borowicz, A., Forrest, S., Witharana, C., Hart, T. and Lynch, H. J. 2020. A global population assessment of the chinstrap penguin (*Pygoscelis antarctica*). *Scientific Reports* 10(1): 19474.
- Testa, G., Neira, S., Giesecke, R., and Piñones, A. 2022. Projecting environmental and krill fishery impacts on the Antarctic Peninsula food web in 2100. *Progress in Oceanography* 206:102862.
- Thiemann, G.W., Iverson, S.J., and Stirling, I. 2008. Polar bear diets and Arctic marine food webs: insights from fatty acid analysis. *Ecological Monographs* 78: 591–613.
- Thomas, P.O., Reeves, R.R., and Brownell Jr., R.L. 2016. Status of the world's baleen whales. *Marine Mammal Science* 32(2):682–734.
- Thompson, T. 2021. 'Arctic sea ice hits 2021 minimum.' *Nature News*, 29 September. <https://www.nature.com/articles/d41586-021-02649-6>.
- Tomaselli, M., Henri, D., Pangnirtung Hunters and Trappers Organization, et al. 2022. Nunavut Inuit Qaujimajatuqangit on the health of the Davis Strait polar bear population. Final project report, Government of Nunavut.
- Trathan, P.N., Wienecke, B., Barbraud, C., et al., including Fretwell, P.T. 2020. The emperor penguin - Vulnerable to projected rates of warming and sea ice loss. *Biological Conservation* 241:108216.
- Turner, J. and Comiso, J. 2017. Solve Antarctica's sea-ice puzzle. *Nature* 547:275–277.
- Turner, J. and Overland, J. 2009. Contrasting climate change in the two polar regions. *Polar Research* 28(2):146–164.
- Turner, J., Bracegirdle, T.J., Phillips, T. et al. 2013. An initial assessment of Antarctic sea ice extent in the CMIP5 models. *Journal of Climate* 26(5):1473–1484.
- US Fish and Wildlife Service (USFWS). 2008. Determination of threatened status for the polar bear (*Ursus maritimus*) throughout its range. *Federal Register* 73: 28212–28303.
- US Fish and Wildlife Service (USFWS). 2016. Polar bear (*Ursus maritimus*) conservation management plan, final. USFWS, Region 7, Anchorage, Alaska.
- USFWS. 2017a. 'After Comprehensive Review, Service Determines Pacific Walrus Does Not Require Endangered Species Act Protection.' *US Fish and Wildlife Service Press Release*, 4 October.
- USFWS. 2017b. 'Endangered and threatened wildlife and plants; 12-month findings on petitions to list 25 species as endangered or threatened species.' *Federal Register* 82(192):46618–46645.
- USFWS 2022a. 'US Fish and Wildlife Service Provides Endangered Species Act Protections for Emperor Penguin.' US Fish and Wildlife Service Press Release, 25 October. <https://www.fws.gov/press-release/2022-10/emperor-penguin-gets-endangered-species-act-protections>.
- USFWS 2022b. 'Endangered and Threatened Wildlife and Plants; Threatened Species Status for Emperor Penguin With Section 4(d) Rule.' *Federal Register* 87(206):64700–64720.
- US Geological Survey (USGS). 2007. Executive Summary, USGS Science Strategy to Support U.S. Fish and Wildlife Service Polar Bear Listing Decision. Administrative Report, US Geological Survey. Reston, Virginia.
- Van Waerebeek, K., Leaper, R., Baker, A. et al. 2010. Odontocetes of the Southern Ocean sanctuary. *Journal of Cetacean Research Management* 11(3):315–346.
- Verma, J.T. 2022. 'This fishing captain is combining Inuit knowledge with scientific expertise to fight climate change in the Far North.' *The Globe and Mail* (Canada), 24 April. <https://www.theglobeandmail.com/canada/article-inuit-climate-change-ice-scientist/>.
- Vibe, C. 1965. The polar bear in Greenland. In: Proceedings of the First International Scientific Meeting on the Polar Bear. Fairbanks, Alaska 6–10 September 1965, *IUCN Polar Bear Specialist Group. University of Alaska International Conference Proceedings Series, No. 1*. pp. 17–25. Washington, DC.

- Vibe 1967. Arctic Animals in Relation to Climatic Fluctuations. *Meddelelser om Grønland* 170(5). C. A. Reitzels Forlag, Copenhagen.
- Wang, M. and Overland, J.E. 2009. A sea ice free summer Arctic within 30 years? *Geophysical Research Letters* 36:L07502. <https://doi.org/10.1029/2009GL037820>.
- WhalesOnline 2016. 'I seem to think I saw a killer whale in Tadoussac this summer; is this possible?' *WhalesOnline Q&A*, 8 November. <https://baleinesendirect.org/en/i-seem-to-think-i-saw-a-killer-whale-in-tadoussac-this-summer-is-this-possible/>.
- Wiig, Ø., Amstrup, S., Atwood, T., et al. 2015. *Ursus maritimus*. The IUCN Red List of Threatened Species 2015: e.T22823A14871490. Available from <http://www.iucnredlist.org/details/22823/0> [accessed Nov. 28, 2015].
- Wiig, Ø., Atkinson, S.N., Born, E.W., et al. 2021. An on-ice aerial survey of the Kane Basin polar bear (*Ursus maritimus*) subpopulation. *Polar Biology* 45:89–100. <https://doi.org/10.1007/s00300-021-02974-6>.
- Wilder, J.M., Vongraven, D., Atwood, T., et al. 2017. Polar bear attacks on humans: implications of a changing climate. *Wildlife Society Bulletin* 41(3):537–547.
- Wilson, B., Crockford, S.J., Johnson, J.W., et al. 2011. Genetic and archaeological evidence of a breeding population of formerly endangered Aleutian cackling goose, *Branta hutchinsii leucopareia*, on Adak Island in the central Aleutians, Alaska. *Canadian Journal of Zoology* 89:732–743.
- Younger, J.L., Emmerson, L.M. and Miller, K.J. 2015. The influence of historical climate changes on Southern Ocean marine predator populations: a comparative analysis. *Global Change Biology* 22(2):474–493.
- Yuan, N., Ding, M., Ludescher, J. and Bunde, A. 2017. Increase of the Antarctic Sea Ice Extent is highly significant only in the Ross Sea. *Scientific Reports* 7:41096.
- Zane, L., Marcato, S., Bargelloni, L. et al. 2006. Demographic history and population structure of the Antarctic silverfish *Pleuragramma antarcticum*. *Molecular Ecology* 15(14):4499–4511.

References

1. The author has published original research results in related fields (most recently: Crockford 2018a; Leathlobhair et al. 2018; McKechnie et al. 2020), research on evolutionary theory that includes polar bears (Crockford 2003, 2004, 2006), research on evolutionary theory that includes geological and atmospheric processes (Crockford 2009), and reviews and synthesis reports on Arctic climate and seals (Crockford 2008, 2012, 2022b; Crockford and Frederick 2007, 2011).
2. <https://susancrockford.com>, all available through *Amazon*.
3. Crockford 2017, 2022b.
4. Crockford 2012; 2014a, b, Crockford 2015a, b, Crockford 2017; Crockford 2018b-e; Crockford 2019a-f; Crockford 2020a,b; 2021; 2022a; 2023.
5. Meier et al. 2022; Omrani et al. 2022.
6. Cornish et al. 2022; Lang et al. 2017; Meier and Stewart 2019; Meier et al. 2022.
7. Shiozaki et al. 2022.
8. Andersen et al. 2021; Ardyna et al. 2020a, b; Bengtsson et al. 2020, 2021; Crawford et al. 2015; Crockford 2022b, Frey et al. 2022; Lewis et al. 2020; Rode et al. 2021; but see Farmer et al. 2021.
9. Meier et al. 2022; Thompson 2021; S.J. Crockford, <https://polarbearsience.com/2021/10/20/most-chukchi-sea-ice-in-20-years-means-no-walrus-feasts-for-polar-bears-at-famous-russian-cliffs/>.
10. Molnár et al. 2010, 2020; see also Hausfather and Peters 2020.
11. Castro de la Guardia et al. 2017; A.E. Derocher tweet 18 June 2022: ('First reports of a polar bear near Churchill are coming in...None of our tracked bears are ashore yet'), <https://twitter.com/AEDerocher/status/1538301855159046144>.
A.E. Derocher tweet 28 July 2022: ('It's almost like the 1980s! This should be great for the bears'), <https://twitter.com/AEDerocher/status/1552842479916892163>; A.E. Derocher tweet 22 August 2022: ('The last polar bear (bear E) is now ashore'), <https://twitter.com/AEDerocher/status/1297193992891768833>.
12. S.J. Crockford, <https://polarbearsience.com/2022/07/19/some-of-the-first-polar-bears-onshore-in-western-hudson-bay-are-in-excellent-condition/> and <https://polarbearsience.com/2022/08/31/video-fat-polar-bears-on-the-shore-of-hudson-bay-near-churchill-one-catches-a-goose/>.
13. Castro de la Guardia et al. 2017; S.J. Crockford, <https://polarbearsience.com/2022/11/17/hudson-bay-sea-ice-freeze-up-in-2022-like-the-1980s-for-the-5th-time-since-2015/>.
14. Castro de la Guardia et al. 2017; Kaufman 2019.
15. Miller et al. 2022; Freeze-up dates: 2020, Nov. 9; 2019, Nov 16; 2018, Nov. 15; 2017, Nov. 15; 2016, Dec. 9; vs. average date for the 1980s of Nov. 16 determined by Castro de la Guardia et al. 2017.
16. Gill 2022; Jacobo 2022.
17. Aars et al. 2006; Crockford 2017; Regehr et al. 2016; Wiig et al. 2015; USFWS 2008; USFWS 2016.
18. COSEWIC 2018; see also <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/polar-bear-2018.html>.
19. PBSG 2021.
20. Bromaghin et al. 2021; Conn et al. 2021; Crockford 2019b, 2022b; Durner et al. 2018; Dyck et al. 2021a; Matishov et al. 2014; PBSG 2021; Regehr et al. 2016, 2018; Wiig et al. 2015.
21. It has been argued (Crockford 2017, 2019b) that a plausible and scientifically defensible 'best-guess' estimate at 2018, extrapolated from 'known' to 'unknown' subpopulations within sea ice ecoregions, would be about 39,000 (range 26,000-58,000), although a more pessimistic best-guess (Hamilton and Derocher 2019) based on a greater variety of ecosystem traits (including

- prey diversity and sea ice cover) came out much lower, at 23,315 (range 15,972-31,212); see also Bromaghin et al. 2021; Crockford 2021; Conn et al. 2021; Dyck et al. 2021a,b, 2023; COSEWIC 2018 ('Total Abundance' section); <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/polar-bear-2018.html>.
22. Amstrup et al. 2007; Crockford 2017, 2019b, 2020b ('The problem of statistical confidence'); USGS 2007.
 23. Meier 2019; Meier et al. 2022; Stroeve et al. 2007, 2014; Wang and Overland 2009.
 24. Crockford 2021; see also Bromaghin et al. 2021; Wiig et al. 2021.
 25. Dyck 2021b; see Atkinson et al. 2021 regarding the genetic sampling method used.
 26. Dyck et al. 2021a, Dyck et al. 2023; Government of Canada 2022; Tomaselli et al. 2022; Snook 2022.
 27. Crockford 2023; Dickie 2022b; Naishadham 2022; Agencie-France Presse 2022.
 28. Dyck et al. 2021b; George 2022; Tomaselli et al. 2022.
 29. Norwegian Polar Institute 2022a; see also NPI 2022b, NPI 2022c; Doran et al. 2020; Lippold et al. 2019.
 30. Jacobo 2022; Wilder et al. 2017.
 31. Environment and Climate Change Canada 2021; George 2022; Tomaselli et al. 2022.
 32. Smith 2022; Smith et al. 2022; but see Dyck et al. 2021b.
 33. Crockford 2019b, 2022a.
 34. Doran et al. 2020; Haavik 2022; Hamamdjian and Jones 2022; Lippold et al. 2019; Miller et al. 2022; Molnar et al. 2020; Richardson 2019; Rode et al. 2014, 2018, 2021; Stirling and Derocher 2012.
 35. Brown et al. 2018; Thiemann et al. 2008.
 36. e.g. Calvert and Stirling 1990.
 37. Herreman and Peacock 2013; Lillie et al. 2019; Miller et al. 2006; Miller et al. 2015; Rogers et al. 2015; See also <http://www.dailymail.co.uk/news/article-5110801/Polar-bears-scramble-mountain-feast-whale.html>.
 38. USFWS 2017a, 2017b; see also Kovacs 2016.
 39. Fischbach et al. 2022.
 40. Holt 2019; Chakilev and Kochnev 2019.
 41. Beatty et al. 2019; MacCracken et al. 2017:25.
 42. Norwegian Polar Institute 2022d.
 43. NAMMCO 2021.
 44. Environment and Climate Change Canada 2021; Martinez-Levasseur et al. 2021.
 45. Tomaselli et al. 2022.
 46. Tomaselli et al. 2022:63.
 47. Bilby 2022; Gamillo 2022.
 48. George et al. 2020.
 49. CBC News 2021; Halliday et al. 2022; WhalesOnline 2016.
 50. Gamillo 2022; Innes et al. 2002; Smith and Martin 1994; COSEWIC 2020 beluga population assessments: Eastern High Arctic/Baffin Bay, 'Special Concern'; Cumberland Sound, 'Endangered'; Ungava Bay, 'Endangered'; W Hudson Bay, 'Not at Risk'; E Hudson Bay, 'Threatened'; James Bay, 'Not at Risk'; DFO 2022 beluga population assessment St. Lawrence Estuary, 'Endangered'.
 51. Pearce et al. 2022.
 52. Wilson et al. 2011.
 53. Kaler et al. 2022.
 54. Doherty 2016.

55. Campana et al. 2020; Dunmall et al. 2022; Siddon 2021; Snoeijs-Leijonmalm et al. 2022; Verma 2022.
56. Aars et al. 2017; Benson and Trites 2002; Fedewa et al. 2020; Heide-Jørgensen et al. 2022; Mangerud and Svendsen 2018; Ogilvie et al. 2000; Ramirez 2022; Stirling and Kiliaan 1980; Vibe 1965; Vibe 1967.
57. Scott 2020.
58. Colucci 2022; Comiso et al. 2017; Fogt et al. 2022; NSIDC 2023; Yuan et al. 2017.
59. Chinchar 2021; NSIDC 2021, 2022.
60. NSIDC 2014.
61. Christie et al. 2022; Collins 2022.
62. Blanchard-Wrigglesworth et al. 2021, 2022; Turner and Comiso 2017; Turner and Overland 2009; Turner et al. 2013.
63. Rackow et al. 2022.
64. e.g. Jenouvrier et al. 2020; Trathan et al. 2020 (emperor penguin); Krüger et al. 2020; McBride et al. 2021; Piñones and Fedorov 2016 (krill); Strycker et al. 2020 (chinstrap penguin).
65. Arce et al. 2022; Pinkerton et al. 2021; Smith Jr. 2022.
66. Siegel and Watkins 2016.
67. Brooks et al. 2022; De Augustinis 2022a; Dickie 2022a; Krüger et al. 2020; Lowther et al. 2020; Meyer et al. 2020; Antarctic krill fishery is managed by the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR).
68. Durban et al. 2017; Durban et al. 2021; Foote et al. 2019; Pitman 2011; Pitman et al. 2007.
69. Van Waerebeek et al. 2010.
70. Durban et al. 2017; Fearnbach et al. 2021.
71. Fearnbach et al. 2019.
72. Pitman 2011; Van Waerebeek et al. 2010.
73. Pitman 2011.
74. Australian Antarctic Program 2020.
75. Cooke et al. 2018; Herr et al. 2022; Perrin and Brownell Jr. 2009; Risch et al. 2019.
76. Cooke 2018b; Herr et al. 2022; IWC 2018b; Ryan 2022.
77. Bestley et al. 2019; IWC 2016.
78. Cooke and Zerbini 2018.
79. Australian Antarctic Program 2022; Cooke 2018a; IWC 2018a; Thomas et al. 2016.
80. Arce et al. 2022; Bailleau et al. 2007; Bornemann et al. 2000; Chua et al. 2022; Labrousse et al. 2018; March et al. 2021.
81. Hückstädt 2015a; LaRue et al. 2021; Marine Mammal Science 2023a.
82. Hückstädt 2015a.
83. Bengtsson 2009; Hückstädt 2015b.
84. Hückstädt 2015c.
85. Hückstädt 2015d.
86. BirdLife International 2020e, f, g.
87. Younger et al. 2015.
88. BirdLife International 2020d.
89. BirdLife International 2020b; Southwell 2017.
90. BirdLife International 2020c; Strycker and Lynch 2022.
91. Stryker 2020; Strycker et al. 2020.
92. Amos 2019, Amos 2020; Fretwell et al. 2012; Fretwell and Trathan 2019, 2020.
93. BirdLife International 2018; Trathan et al. 2020.

94. Crockford 2019b; Hausfather and Peters 2020; Jenouvrier et al. 2009; Jenouvrier et al. 2020.
95. Blanchard-Wrigglesworth et al. 2022; Rackow et al. 2022; Turner and Comiso 2017; Turner and Overland 2009; Turner et al. 2013.
96. BirdLife International 2020a.
97. Federal Register 2022; USFWS 2022a, b.
98. e.g. Carneiro et al. 2022; Collins et al. 2021.
99. Brooks et al. 2022; De Augustinis 2022b.
100. Zane et al. 2006.

Review process

GWPF publishes papers in a number of different formats, with a different review process pertaining to each.

- Our flagship long-form GWPF Reports are all reviewed by our Academic Advisory Council.
- GWPF Briefings and Notes are shorter documents and are reviewed internally and/or externally as required.

Part of the function of the review process is to ensure that any materials published by the GWPF are of a proper academic standard, and will serve the GWPF's educational purpose. As a charity, we recognise that educational material should provide any reader the opportunity to understand, and explore different perspectives on a subject.

This means that, for most publications, we also invite an external review from a party who we would expect to take a different view to the publication's author. We offer to publish any substantive comments alongside the main paper, provided we are satisfied they will enhance the educational experience of the reader. In this way, we hope to encourage open and active debate on the important areas in which we work.

This enhanced review process for GWPF papers is intended to take the content and analysis beyond a typical review for an academic journal:

- More potential reviewers can be involved
- The number of substantive comments will typically exceed journal peer review, and
- The identity of the author is known to the potential reviewers.

As an organisation whose publications are sometimes the subject of assertive or careless criticism, this review process is intended to enhance the educational experience for all readers, allowing points to be made and considered in context and observing the standards required for an informed and informative debate. We therefore expect all parties involved to treat the reviews with the utmost seriousness.

Final responsibility for publication rests with the Chairman of the Trustees and the GWPF Director. But in every case, the views expressed are those of the author. GWPF has never had any corporate position beyond that dictated by its educational objectives.

About the Global Warming Policy Foundation

People are naturally concerned about the environment, and want to see policies that protect it, while enhancing human wellbeing; policies that don't hurt, but help.

The Global Warming Policy Foundation (GWPF) is committed to the search for practical policies. Our aim is to raise standards in learning and understanding through rigorous research and analysis, to help inform a balanced debate amongst the interested public and decision-makers. We aim to create an educational platform on which common ground can be established, helping to overcome polarisation and partisanship. We aim to promote a culture of debate, respect, and a hunger for knowledge.

Views expressed in the publications of the Global Warming Policy Foundation are those of the authors, not those of the GWPF, its trustees, its Academic Advisory Council members or its directors.

THE GLOBAL WARMING POLICY FOUNDATION

Director

Benny Peiser

Honorary President

Lord Lawson

BOARD OF TRUSTEES

Dr Jerome Booth (Chairman)

The Hon. Tony Abbott

Lord Frost

Kathy Gyngell

Professor Michael Kelly FRS

Terence Mordaunt

Graham Stringer MP

Professor Fritz Vahrenholt

ACADEMIC ADVISORY COUNCIL

Professor Christopher Essex (Chairman)

Professor Wade Allison

Professor Anthony Barrett

Professor J. Ray Bates

Sir Ian Byatt

Dr John Constable

Professor Vincent Courtillot

Professor John Dewey

Professor Peter Dobson

Professor Peter Edwards FRS

Professor Samuel Furfari

Christian Gerondeau

Professor Larry Gould

Professor William Happer

Professor Ole Humlum

Professor Gautam Kalghatgi

Professor Terence Kealey

Bill Kininmonth

Brian Leyland

Professor Richard Lindzen

Professor Ross McKittrick

Professor Robert Mendelsohn

Professor Garth Paltridge

Professor Ian Plimer

Professor Gwythian Prins

Professor Paul Reiter

Professor Peter Ridd

Dr Matt Ridley

Sir Alan Rudge

Professor Nir Shaviv

Professor Henrik Svensmark

Dr David Whitehouse

RECENT GWPF BRIEFINGS

28	Susan Crockford	20 Good Reasons not to Worry About Polar Bears: Update
29	Mikko Paunio	Sacrificing the Poor: The Lancet on 'pollution'
30	Mikko Paunio	Kicking Away the Energy Ladder
31	Bill Gray	Flaws in Applying Greenhouse Warming to Climate Variability
32	Mikko Paunio	Save the Oceans: Stop Recycling Plastic
33	Andy Dawson	Small Modular Nuclear: Crushed at Birth
34	Andrew Montford	Quakes, Pollution and Flaming Faucets
35	Paul Homewood	DEFRA vs Met Office: Factchecking the State of the UK Climate
36	J. Ray Bates	Deficiencies in the IPCC's Special Report on 1.5 Degrees
37	Paul Homewood	Tropical Hurricanes in the Age of Global Warming
38	Mikko Paunio	The Health Benefits of Ignoring the IPCC
39	Jack Ponton	Grid-scale Storage: Can it Solve the Intermittency Problem?
40	Robert Lyman	Carbon Taxation: The Canadian Experience
41	Rémy Prud'homme	La Transition Énergétique: Useless, Costly, Unfair
42	Judith Curry	Recovery, Resilience, Readiness: Contending with Natural Disasters
43	Paul Homewood	Plus Ça Change: The UK Climate in 2018
44	David Whitehouse	Cold Water: The Oceans and Climate Change
45	Crockford and Laframboise	The Defenestration of Dr Crockford
46	Paul Homewood	Britain's Weather in 2019: More of the Same, Again
47	John Constable	The Brink of Darkness: Britain's Fragile Grid
48	Mike Travers	The Hidden Cost of Net Zero: Rewiring the UK
49	Martin Livermore	Greenhouse Gas Emissions: The Global Picture
50	Paul Homewood	The US Climate in 2019
51	Patricia Adams	The Red and the Green: China's Useful Idiots
52	Andrew Montford	Offshore Wind: Cost Predictions and Cost Outcomes
53	Tim Worstall	A Saviour Spurned: How Fracking Saved us from Global Warming
54	Jun Arima	Eco-fundamentalism as Grist for China's Mill
55	Gautam Kalghatgi	Scoping Net Zero
56	Andrew Montford	Survival of the Richest: Smart Homes and Energy Rationing
57	Donna Laframboise	The Hounding of Roger Pielke Jr
58	Patricia Adams	China's Energy Dream
59	Andrew Montford	The Rising Cost of Onshore Wind
60	Paul Homewood	The UK's Weather in 2020-21
61	Francis Menton	The Energy Storage Conundrum
62	Paul Homewood	The 2022 Hurricane Season
63	Susan Crockford	The Polar Wildlife Report

For further information about the Global Warming Policy Foundation, please visit our website at www.thegwpf.org.

The GWPF is a registered charity, number 1131448.

